

P-cadherin (A-10): sc-74545

BACKGROUND

Cadherins comprise a family of Ca^{2+} -dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. The classical cadherins, E-, N- and P-cadherin, consist of large extracellular domains characterized by a series of five homologous NH_2 terminal repeats. The most distal of these cadherins is thought to be responsible for binding specificity, transmembrane domains and carboxy-terminal intracellular domains. The relatively short intracellular domains interact with a variety of cytoplasmic proteins, such as β -catenin, to regulate cadherin function. Members of this family of adhesion proteins include rat cadherin K (and its human homolog, cadherin-6), R-cadherin, B-cadherin, E/P cadherin and cadherin-5.

REFERENCES

1. Takeichi, M. 1988. The cadherins: cell-cell adhesion molecules controlling animal morphogenesis. *Development* 102: 639-655.
2. Hatta, M., et al. 1991. Genomic organization and chromosomal mapping of the mouse P-cadherin gene. *Nucleic Acids Res.* 19: 4437-4441.
3. Koch, P.J., et al. 1994. Desmosomal cadherins: another growing multigene family of adhesion molecules. *Curr. Opin. Cell Biol.* 6: 682-687.
4. Ranscht, B. 1994. Cadherins and catenins: interactions and functions in embryonic development. *Curr. Opin. Cell Biol.* 6: 740-746.
5. Hinck, L., et al. 1994. Dynamics of cadherin/catenin complex formation: novel protein interactions and pathways of complex assembly. *J. Cell Biol.* 125: 1327-1340.

CHROMOSOMAL LOCATION

Genetic locus: CDH3 (human) mapping to 16q22.1.

SOURCE

P-cadherin (A-10) is a mouse monoclonal antibody raised against amino acids 550-654 mapping within an extracellular domain of P-cadherin of human origin.

PRODUCT

Each vial contains 200 μg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

P-cadherin (A-10) is available conjugated to agarose (sc-74545 AC), 500 μg /0.25 ml agarose in 1 ml, for IP; to HRP (sc-74545 HRP), 200 μg /ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-74545 PE), fluorescein (sc-74545 FITC), Alexa Fluor[®] 488 (sc-74545 AF488), Alexa Fluor[®] 546 (sc-74545 AF546), Alexa Fluor[®] 594 (sc-74545 AF594) or Alexa Fluor[®] 647 (sc-74545 AF647), 200 μg /ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-74545 AF680) or Alexa Fluor[®] 790 (sc-74545 AF790), 200 μg /ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS


P-cadherin (A-10) is recommended for detection of P-cadherin of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for P-cadherin siRNA (h): sc-29420, P-cadherin shRNA Plasmid (h): sc-29420-SH and P-cadherin shRNA (h) Lentiviral Particles: sc-29420-V.


Molecular Weight of P-cadherin: 118 kDa.

Positive Controls: Caco-2 cell lysate: sc-2262, PC-3 cell lysate: sc-2220 or A-431 whole cell lysate: sc-2201.

DATA



P-cadherin (A-10): sc-74545. Western blot analysis of P-cadherin expression in A-431 (A) and Caco-2 (B) whole cell lysates.



P-cadherin (A-10): sc-74545. Immunoperoxidase staining of formalin fixed, paraffin-embedded human tonsil tissue showing membrane and cytoplasmic staining of squamous epithelial cells.

SELECT PRODUCT CITATIONS

1. Wakamatsu, K., et al. 2022. Metabolites and biomarker compounds of neurodegenerative diseases in cerebrospinal fluid. *Metabolites* 12: 343.
2. Takebayashi, G., et al. 2023. E-cadherin is expressed in epithelial cells of the choroid plexus in human and mouse brains. *Curr. Issues Mol. Biol.* 45: 7813-7826.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.