

GluRS (S-20): sc-74718

BACKGROUND

The fidelity of protein synthesis requires efficient discrimination of amino acid substrates by aminoacyl-tRNA synthetases. Aminoacyl-tRNA synthetases function to catalyze the aminoacylation of tRNAs by their corresponding amino acids, thus linking amino acids with tRNA-contained nucleotide triplets. GluRS (glutamyl-tRNA synthetase 2), also known as EARS2 or MSE1, is a 523 amino acid protein that localizes to the mitochondrial matrix and belongs to the class I aminoacyl-tRNA synthetase family. Participating in protein biosynthesis, GluRS functions to catalyze the ATP-dependent attachment of glutamate to tRNA(Glu), a two-step reaction that involves the ATP-dependent activation of glutamate to form Glu-AMP and the subsequent transfer of the glutamate residue to tRNA(Glu).

REFERENCES

1. Frugier, M., Söll, D., Giege, R. and Florentz, C. 1994. Identity switches between tRNAs aminoacylated by class I glutamyl- and class II aspartyl-tRNA synthetases. *Biochemistry* 33: 9912-9921.
2. Freist, W., Gauss, D.H., Söll, D. and Lapointe, J. 1997. Glutamyl-tRNA synthetase. *Biol. Chem.* 378: 1313-1329.
3. Quevillon, S., Robinson, J.C., Berthonneau, E., Siatecka, M. and Mirande, M. 1999. Macromolecular assemblage of aminoacyl-tRNA synthetases: identification of protein-protein interactions and characterization of a core protein. *J. Mol. Biol.* 285: 183-195.
4. Robinson, J.C., Kerjan, P. and Mirande, M. 2000. Macromolecular assemblage of aminoacyl-tRNA synthetases: quantitative analysis of protein-protein interactions and mechanism of complex assembly. *J. Mol. Biol.* 304: 983-994.
5. Han, J.M., Kim, J.Y. and Kim, S. 2003. Molecular network and functional implications of macromolecular tRNA synthetase complex. *Biochem. Biophys. Res. Commun.* 303: 985-993.
6. Lee, S.W., Cho, B.H., Park, S.G. and Kim, S. 2004. Aminoacyl-tRNA synthetase complexes: beyond translation. *J. Cell Sci.* 117: 3725-3734.
7. Bonnefond, L., Fender, A., Rudinger-Thirion, J., Giege, R., Florentz, C. and Sissler, M. 2005. Toward the full set of human mitochondrial aminoacyl-tRNA synthetases: characterization of AspRS and TyrRS. *Biochemistry* 44: 4805-4816.
8. Sethi, A., Eargle, J., Black, A.A. and Luthey-Schulten, Z. 2009. Dynamical networks in tRNA:protein complexes. *Proc. Natl. Acad. Sci. USA* 106: 6620-6625.

CHROMOSOMAL LOCATION

Genetic locus: EARS2 (human) mapping to 16p12.1; Ears2 (mouse) mapping to 7 F3.

SOURCE

GluRS (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of GluRS of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-74718 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GluRS (S-20) is recommended for detection of GluRS of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GluRS (S-20) is also recommended for detection of GluRS in additional species, including equine, bovine, porcine and avian.

Suitable for use as control antibody for GluRS siRNA (h): sc-75146, GluRS siRNA (m): sc-75147, GluRS shRNA Plasmid (h): sc-75146-SH, GluRS shRNA Plasmid (m): sc-75147-SH, GluRS shRNA (h) Lentiviral Particles: sc-75146-V and GluRS shRNA (m) Lentiviral Particles: sc-75147-V.

Molecular Weight of GluRS: 59 kDa.

Positive Controls: mouse testis extract: sc-2405.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.