

opticin (L-15): sc-74773

BACKGROUND

The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic α/β horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. Opticin, also known as OPTC or OPT, is a 332 amino acid protein that is secreted into the extracellular matrix and contains seven LRR repeats. Expressed in skin and fetal liver and also present in ocular tissues, such as the retina, cornea and iris, opticin functions to bind collagen fibrils and, via this interaction, is thought to regulate fibril spacing, morphology and organization. Opticin is subject to posttranslational O-glycosylation and may be associated with the pathogenesis of primary open angle glaucoma.

REFERENCES

1. Reardon, A.J., Le Goff, M., Briggs, M.D., McLeod, D., Sheehan, J.K., Thornton, D.J. and Bishop, P.N. 2000. Identification in vitreous and molecular cloning of opticin, a novel member of the family of leucine-rich repeat proteins of the extracellular matrix. *J. Biol. Chem.* 275: 2123-2129.
2. Hobby, P., Wyatt, M.K., Gan, W., Bernstein, S., Tomarev, S., Slingsby, C. and Wistow, G. 2000. Cloning, modeling, and chromosomal localization for a small leucine-rich repeat proteoglycan (SLRP) family member expressed in human eye. *Mol. Vis.* 6: 72-78.
3. Friedman, J.S., Faucher, M., Hiscott, P., Biron, V.L., Malenfant, M., Turcotte, P., Raymond, V. and Walter, M.A. 2002. Protein localization in the human eye and genetic screen of opticin. *Hum. Mol. Genet.* 11: 1333-1342.
4. Le Goff, M.M., Hindson, V.J., Jowitt, T.A., Scott, P.G. and Bishop, P.N. 2003. Characterization of opticin and evidence of stable dimerization in solution. *J. Biol. Chem.* 278: 45280-45287.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 605127. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Ramesh, S., Bonshek, R.E. and Bishop, P.N. 2004. Immunolocalisation of opticin in the human eye. *Br. J. Ophthalmol.* 88: 697-702.
7. Acharya, M., Mookherjee, S., Bhattacharjee, A., Thakur, S.K., Bandyopadhyay, A.K., Sen, A., Chakrabarti, S. and Ray, K. 2007. Evaluation of the OPTC gene in primary open angle glaucoma: functional significance of a silent change. *BMC Mol. Biol.* 8: 21.
8. Kumar, A., Basavaraj, M.G., Gupta, S.K., Qamar, I., Ali, A.M., Bajaj, V., Ramesh, T.K., Prakash, D.R., Shetty, J.S. and Dorairaj, S.K. 2007. Role of CYP1B1, MYOC, OPTN, and OPTC genes in adult-onset primary open-angle glaucoma: predominance of CYP1B1 mutations in Indian patients. *Mol. Vis.* 13: 667-676.

CHROMOSOMAL LOCATION

Genetic locus: OPTC (human) mapping to 1q32.1; Optc (mouse) mapping to 1 E4.

SOURCE

opticin (L-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of opticin of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-74773 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

opticin (L-15) is recommended for detection of opticin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for opticin siRNA (h): sc-75996, opticin siRNA (m): sc-75997, opticin shRNA Plasmid (h): sc-75996-SH, opticin shRNA Plasmid (m): sc-75997-SH, opticin shRNA (h) Lentiviral Particles: sc-75996-V and opticin shRNA (m) Lentiviral Particles: sc-75997-V.

Molecular Weight of opticin: 45 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **opticin (F-2): sc-271306**, our highly recommended monoclonal alternative to opticin (L-15).