

CD72 (H-7): sc-7483

BACKGROUND

CD5 has been identified as a transmembrane glycoprotein that is expressed on 70% of normal peripheral blood lymphocytes and on virtually all T lymphocytes in thymus and peripheral blood. Activation of T cells through the T cell receptor (TCR) results in tyrosine phosphorylation of CD5, and the absence of CD5 renders T cells hyper-responsive to TCR-mediated activation. CD5 associates with the TCR/CD3 ζ chain, and with the Src family kinase, Lck p56. The C-type lectin superfamily member CD72 is a cell surface negative regulator of B cell activation from the pro-B through the mature B cell stage. CD72 serves as a receptor for CD5. The ability of lymphocytes to respond to antigenic or mitogenic stimulation utilizes both positive and negative regulatory proteins that influence the threshold for responsiveness. The human CD72 gene maps to chromosome 9p13.3 and encodes a transmembrane glycoprotein that contains an immunoreceptor tyrosine-based inhibition motif (ITIM). Upon tyrosine phosphorylation, the CD72 ITIM recruits SH2-containing phosphatases such as SHP-1, resulting in downregulation of cell activation. CD72^{-/-} mice contain hyperproliferative B cells.

REFERENCES

- Knapp, W., et al, eds. 1989. Leucocyte Typing IV. New York: Oxford University Press.
- Schlossman, S., et al, eds. 1995. Leucocyte Typing V. Oxford: Oxford University Press.
- Ying, H., et al. 1998. Regulation of mouse CD72 gene expression during B lymphocyte development. *J. Immunol.* 161: 4760-4767.
- Wu, Y., et al. 1998. The B cell transmembrane protein CD72 binds to and is an *in vivo* substrate of the protein tyrosine phosphatase SHP-1. *Curr. Biol.* 8: 1009-1017.
- Pan, C., et al. 1999. CD72-deficient mice reveal nonredundant roles of CD72 in B cell development and activation. *Immunity* 11: 495-506.
- Parnes, J.R., et al. 2000. CD72, a negative regulator of B cell responsiveness. *Immunol. Rev.* 176: 75-85.
- Wu, H.J., et al. 2002. Positive and negative roles of CD72 in B cell function. *Immunol. Res.* 25: 155-166.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 107272. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: CD72 (human) mapping to 9p13.3; Cd72 (mouse) mapping to 4 B1.

SOURCE

CD72 (H-7) is a mouse monoclonal antibody raised against amino acids 1-96 mapping at the N-terminus of CD72 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CD72 (H-7) is recommended for detection of CD72 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for CD72 siRNA (h): sc-37250, CD72 siRNA (m): sc-35022, CD72 shRNA Plasmid (h): sc-37250-SH, CD72 shRNA Plasmid (m): sc-35022-SH, CD72 shRNA (h) Lentiviral Particles: sc-37250-V and CD72 shRNA (m) Lentiviral Particles: sc-35022-V.


Molecular Weight of CD72: 45 kDa.

Positive Controls: rat spleen extract: sc-2397, IB4 whole cell lysate: sc-364780 or WEHI-231 whole cell lysate: sc-2213.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



CD72 (H-7): sc-7483. Western blot analysis of CD72 expression in IB4 (A), WEHI-231 (B), Raji (C) and Jurkat (D) whole cell lysates and rat spleen tissue extract (E).



CD72 (H-7): sc-7483. Immunoperoxidase staining of formalin fixed, paraffin-embedded mouse lymphoma tissue showing membrane localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.