

# ACADVL (G-16): sc-74898

## BACKGROUND

ACADVL (acyl-coenzyme A dehydrogenase, very long chain), also known as VLCAD, LCACD or ACAD6, is an inner mitochondrial membrane protein that belongs to the family of acyl-CoA dehydrogenases. Involved in lipid metabolism, ACADVL has catalytic activity toward esters of long chain and very long chain fatty acids and functions in the first step of the fatty acid  $\beta$ -oxidation pathway. Defects in the gene encoding ACADVL are the cause of very long chain acyl-CoA dehydrogenase deficiency (VLCAD deficiency), an autosomal recessive disease that leads to impaired long-chain fatty acid oxidation. VLCAD deficiency can give rise to three different phenotypes: a severe childhood form with a high incidence of mortality and cardiomyopathy; a mild childhood form with hypoketotic hypoglycemia and low mortality rates; and an adult form characterized by myoglobinuria and rhabdomyolysis. Two isoforms of ACADVL exist due to alternative splicing events.

## CHROMOSOMAL LOCATION

Genetic locus: ACADVL (human) mapping to 17p13.1; Acadvl (mouse) mapping to 11 B3.

## SOURCE

ACADVL (G-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ACADVL of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-74898 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

ACADVL (G-16) is recommended for detection of ACADVL of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ACADVL (G-16) is also recommended for detection of ACADVL in additional species, including equine.

Suitable for use as control antibody for ACADVL siRNA (h): sc-72427, ACADVL siRNA (m): sc-72428, ACADVL shRNA Plasmid (h): sc-72427-SH, ACADVL shRNA Plasmid (m): sc-72428-SH, ACADVL shRNA (h) Lentiviral Particles: sc-72427-V and ACADVL shRNA (m) Lentiviral Particles: sc-72428-V.


Molecular Weight of ACADVL: 66 kDa.

Positive Controls: HeLa nuclear extract: sc-2120.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



ACADVL (G-16): sc-74898. Western blot analysis of ACADVL expression in HeLa nuclear extract.

## SELECT PRODUCT CITATIONS

- Dai, D.F., et al. 2012. Mitochondrial proteome remodelling in pressure overload-induced heart failure: the role of mitochondrial oxidative stress. *Cardiovasc. Res.* 93: 79-88.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **ACADVL (H-7): sc-376239** or **ACADVL (D-11): sc-271225**, our highly recommended monoclonal alternatives to ACADVL (G-16).