SANTA CRUZ BIOTECHNOLOGY, INC.

SNAP 25 (N-19): sc-7539



BACKGROUND

Syntaxins were originally thought to be docking proteins, but have now been categorized as anchoring proteins that anchor themselves to the cytoplasmic surfaces of cellular membranes. Syntaxins have been shown to bind to various proteins involved in exocytosis, including VAMPs (vesicle-associated membrane proteins), NSF (N-ethylmaleimide-sensitive factor), SNAP 25, SNAPs (soluble NSF attachment proteins) and Synaptotagmin. VAMPs, also designated synaptobrevins, including VAMP-1 and VAMP-2, and Synaptotagmin, a protein that may function as an inhibitor of exocytosis, are vesicular proteins. SNAPs, including α - and γ -SNAP, are cytoplasmic proteins that bind to a membrane receptor complex composed of VAMP, SNAP 25 and Syntaxin. SNAPs mediate the membrane binding of NSF, which is essential for membrane fusion reactions. An additional protein designated synaptophysin may regulate exocytosis by competing with SNAP 25 and syntaxins for VAMP binding.

CHROMOSOMAL LOCATION

Genetic locus: SNAP25 (human) mapping to 20p12.2; Snap25 (mouse) mapping to 2 F3.

SOURCE

SNAP 25 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of SNAP 25 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-7539 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SNAP 25 (N-19) is recommended for detection of SNAP 25 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SNAP 25 (N-19) is also recommended for detection of SNAP 25 in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for SNAP 25 siRNA (h): sc-36517, SNAP 25 siRNA (m): sc-36516, SNAP 25 shRNA Plasmid (h): sc-36517-SH, SNAP 25 shRNA Plasmid (m): sc-36516-SH, SNAP 25 shRNA (h) Lentiviral Particles: sc-36517-V and SNAP 25 shRNA (m) Lentiviral Particles: sc-36516-V.

Molecular Weight of SNAP 25: 25 kDa.

Positive Controls: SNAP 25 (h): 293 Lysate: sc-111113, SH-SY5Y cell lysate: sc-3812 or PC-12 cell lysate: sc-2250.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA





SNAP 25 (N-19): sc-7539. Western blot analysis of SNAP 25 expression in human SNAP 25 transfected 293: sc-111113 (**A**), non-transfected 293: sc-110760 (**B**) and SH-SY5Y (**C**) whole cell lysates. SNAP 25 (N-19): sc-7539. Immunofluorescence staining of methanol-fixed PC-12 cells showing membrane localization.

SELECT PRODUCT CITATIONS

- Gibbins, I.L., et al. 2003. Heterogeneous expression of SNAP 25 and synaptic vesicle proteins by central and peripheral inputs to sympathetic neurons. J. Comp. Neurol. 459: 25-43.
- Brahmaraju, M., et al. 2004. Spatio-temporal organization of Vam6P and SNAP on mouse spermatozoa and their involvement in sperm-zona pellucida interactions. Biochem. Biophys. Res. Commun. 318: 148-155.
- Morris, J.L., et al. 2005. Most peptide-containing sensory neurons lack proteins for exocytotic release and vesicular transport of glutamate. J. Comp. Neurol. 483: 1-16.
- Kaltenbach, L.S., et al. 2007. Huntingtin interacting proteins are genetic modifiers of neurodegeneration. PLoS Genet. 3: e82.
- Shu, Y., et al. 2008. Phosphorylation of SNAP 25 at Ser187 mediates enhancement of exocytosis by a phorbol ester in INS-1 cells. J. Neurosci. 28: 21-30.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS Satisfation Guaranteed

Try SNAP 25 (SP12): sc-20038 or SNAP 25 (H-1): sc-376713, our highly recommended monoclonal alternatives to SNAP 25 (N-19).