

DcR2 (C-20): sc-7550

BACKGROUND

Tumor necrosis factor (TNF) is a pleiotropic cytokine whose function is mediated by two distinct cell surface receptors, designated TNF-R1 and TNF-R2, which are expressed on most cell types. TNF function is primarily mediated through TNF-R1 signaling. Both TNF-R1 and TNF-R2 belong to the growing TNF receptor superfamily which includes FAS antigen and CD40. TNF-R1 contains a cytoplasmic motif, termed the "death domain" that has been found to be necessary for the transduction of the apoptotic signal. The death domain is also found in several other receptors, including FAS, DR2 (or TRUNDD), DR3 (death receptor 3), DR4, DR5 and DR6. TRUNDD, DR4 and DR5 are receptors for the apoptosis-inducing cytokine TRAIL. Non-death domain-containing receptors, designated decoy receptor (DcR1 or TRID, DcR2 and DcR3), associate with specific ligands and may play a role in cellular resistance to apoptotic stimuli.

REFERENCES

1. Tartaglia, L.A., et al. 1993. A novel domain within the 55 kDa TNF receptor signals cell death. *Cell* 74: 845-853.
2. Smith, C.A., et al. 1994. The TNF receptor superfamily of cellular and viral proteins: activation, costimulation and death. *Cell* 76: 959-962.
3. Nagata, S., et al. 1995. The FAS death factor. *Science* 267: 1449-1456.

CHROMOSOMAL LOCATION

Genetic locus: TNFRSF10D (human) mapping to 8p21.3.

SOURCE

DcR2 (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of DcR2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-7550 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DcR2 (C-20) is recommended for detection of DcR2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DcR2 siRNA (h): sc-35185, DcR2 shRNA Plasmid (h): sc-35185-SH and DcR2 shRNA (h) Lentiviral Particles: sc-35185-V.


Molecular Weight of DcR2: 42 kDa.

Positive Controls: DcR2 (h): 293T Lysate: sc-371573, U-937 cell lysate: sc-2239 or SW480 cell lysate: sc-2219.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



DcR2 (C-20): sc-7550. Western blot analysis of DcR2 expression in non-transfected: sc-117752 (A) and human DcR2 transfected: sc-371573 (B) 293T whole cell lysates.



SDcR2 (C-20): sc-7550. Western blot analysis of human recombinant DcR2 fusion protein.

SELECT PRODUCT CITATIONS

1. Vindrieux, D., et al. 2011. Down-regulation of DcR2 sensitizes androgen-dependent prostate cancer LNCaP cells to TRAIL-induced apoptosis. *Cancer Cell Int.* 11: 42.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **DcR2 (B-P30): sc-65310**, our highly recommended monoclonal alternative to DcR2 (C-20).