

VAMP-1/2/3 (N-20): sc-7579

BACKGROUND

Syntaxins were originally thought to be docking proteins, but have more recently been categorized as anchoring proteins that anchor themselves to the cytoplasmic surfaces of cellular membranes. Syntaxins have been shown to bind to various proteins involved in exocytosis, including VAMPs (vesicle-associated membrane proteins), NSF (N-ethylmaleimide-sensitive factor), SNAP 25 (synaptosomal-associated protein of 25 kDa), SNAPs (soluble NSF attachment proteins) and synaptotagmin. VAMPs, also designated synaptobrevins, including VAMP-1 and VAMP-2, and synaptotagmin, a protein that may function as an inhibitor of exocytosis, are vesicular proteins. SNAPs, including α - and γ -SNAP, are cytoplasmic proteins that bind to a membrane receptor complex composed of VAMP, SNAP 25 and Syntaxin. SNAPs mediate the membrane binding of NSF, which is essential for membrane fusion reactions. An additional protein designated synaptophysin may regulate exocytosis by competing with SNAP 25 and syntaxins for VAMP binding.

REFERENCES

1. Elferink, L.A., et al. 1993. A role for synaptotagmin (p65) in regulated exocytosis. *Cell* 72: 153-159.
2. Bennett, M.K., et al. 1993. The syntaxin family of vesicular transport receptors. *Cell* 74: 863-873.
3. Yamaguchi, K. and Akagawa, K. 1994. Exocytosis relating proteins in the nervous system. *Neurosci. Res.* 20: 289-292.

SOURCE

VAMP-1/2/3 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of VAMP-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-7579 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

VAMP-1/2/3 (N-20) is recommended for detection of VAMP-1, VAMP-2 and VAMP-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

VAMP-1/2/3 (N-20) is also recommended for detection of VAMP-1, VAMP-2 and VAMP-3 in additional species, including equine, canine, bovine, porcine and avian.


Molecular Weight of VAMP-1/2/3: 18 kDa.

Positive Controls: mouse brain extract: sc-2253 or rat brain extract: sc-2392.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



VAMP-1/2/3 (N-20): sc-7579. Western blot analysis of VAMP-1/2/3 expression in mouse brain tissue extract.



VAMP-1/2/3 (N-20): sc-7579. Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing cytoplasmic staining of exocrine glandular cells and Islets of Langerhans.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **VAMP-1/2/3 (F-11): sc-133129** or **VAMP-1/2 (SP10): sc-20039**, our highly recommended monoclonal alternatives to VAMP-1/2/3 (N-20).