

GluR-1 (C-19): sc-7609

BACKGROUND

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration. Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Kainate/AMPA receptors are co-localized with NMDA receptors in many synapses and consist of seven structurally related subunits designated GluR-1 to -7. The kainate/AMPA receptors are primarily responsible for the fast excitatory neuro-transmission by glutamate whereas the NMDA receptors are functionally characterized by a slow kinetic and a high permeability for Ca²⁺ ions. The NMDA receptors consist of five subunits: ϵ 1, 2, 3, 4 and one ζ subunit. The ζ subunit is expressed throughout the brainstem, whereas the four ϵ subunits display limited distribution.

CHROMOSOMAL LOCATION

Genetic locus: GRIA1 (human) mapping to 5q33.2; Gria1 (mouse) mapping to 11 B1.3.

SOURCE

GluR-1 (C-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of GluR-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-7609 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GluR-1 (C-19) is recommended for detection of GluR-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GluR-1 (C-19) is also recommended for detection of GluR-1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for GluR-1 siRNA (h): sc-35485, GluR-1 siRNA (m): sc-35486, GluR-1 shRNA Plasmid (h): sc-35485-SH, GluR-1 shRNA Plasmid (m): sc-35486-SH, GluR-1 shRNA (h) Lentiviral Particles: sc-35485-V and GluR-1 shRNA (m) Lentiviral Particles: sc-35486-V.

Molecular Weight of GluR-1: 106 kDa.

Positive Controls: H4 cell lysate: sc-2408, mouse brain extract: sc-2253 or rat cerebellum extract: sc-2398.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



GluR-1 (C-19): sc-7609. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human brain tumor showing membrane staining.

SELECT PRODUCT CITATIONS

- Farhana, L., et al. 2011. Maximal adamantyl-substituted retinoid-related molecule-induced apoptosis requires NF κ B noncanonical and canonical pathway activation. *Cell Death Differ.* 18: 164-173.
- Yen, Y.H., et al. 2011. A study of the spatial protein organization of the postsynaptic density isolated from porcine cerebral cortex and cerebellum. *Mol. Cell. Proteomics* 10: M110.007138.
- Seese, R.R., et al. 2012. LTP induction translocates cortactin at distant synapses in wild-type but not Fmr1 knock-out mice. *J. Neurosci.* 32: 7403-7413.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **GluR-1 (G-12): sc-55509** or **GluR-1 (E-6): sc-13152**, our highly recommended monoclonal alternatives to GluR-1 (C-19). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **GluR-1 (G-12): sc-55509**.