

ZNF81 siRNA (h): sc-77007

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF81, also known as HFZ20 or MRX45, is a transcriptional regulator belonging to the Krüppel C₂H₂-type zinc-finger protein family. It localizes to the nucleus and contains 12 C₂H₂-type zinc fingers and one KRAB domain. Mutations in the gene encoding ZNF81 are implicated in nonsyndromic X-linked mental retardation (XLMR).

REFERENCES

1. Knight, J.C., Grimaldi, G., Thiesen, H.J., Bech-Hansen, N.T., Fletcher, C.D. and Coleman, M.P. 1994. Clustered organization of Krüppel zinc-finger genes at Xp11.23, flanking a translocation breakpoint at OATL1: a physical map with locus assignments for ZNF21, ZNF41, ZNF81, and ELK1. *Genomics* 21: 180-187.
2. Hagemann, T., Surosky, R., Monaco, A.P., Lehrach, H., Rosen, F.S. and Kwan, S.P. 1994. Physical mapping in a YAC contig of 11 markers on the human X chromosome in Xp11.23. *Genomics* 21: 262-265.
3. Coleman, M.P., Nemeth, A.H., Campbell, L., Raut, C.P., Weissenbach, J. and Davies, K.E. 1994. A 1.8-Mb YAC contig in Xp11.23: identification of CpG islands and physical mapping of CA repeats in a region of high gene density. *Genomics* 21: 337-343.
4. Derry, J.M., Jess, U. and Francke, U. 1996. Cloning and characterization of a novel zinc finger gene in Xp11.2. *Genomics* 30: 361-365.
5. Kleefstra, T., Yntema, H.G., Oudakker, A.R., Banning, M.J., Kalscheuer, V.M., Chelly, J., Moraine, C., Ropers, H.H., Fryns, J.P., Janssen, I.M., Sistermans, E.A., Nillesen, W.N., de Vries, L.B., Hamel, B.C. and van Bokhoven, H. 2004. Zinc finger 81 (ZNF81) mutations associated with X-linked mental retardation. *J. Med. Genet.* 41: 394-399.
6. Martinez-Castellano, F. 2006. Non-specific X-linked mental retardation. *Rev. Neurol.* 42: S77-S83.
7. Lugtenberg, D., Yntema, H.G., Banning, M.J., Oudakker, A.R., Firth, H.V., Willatt, L., Raynaud, M., Kleefstra, T., Fryns, J.P., Ropers, H.H., Chelly, J., Moraine, C., Gecz, J., van Reeuwijk, J., Nabuurs, S.B., de Vries, B.B., et al. 2006. ZNF674: a new kruppel-associated box-containing zinc-finger gene involved in nonsyndromic X-linked mental retardation. *Am. J. Hum. Genet.* 78: 265-278.

CHROMOSOMAL LOCATION

Genetic locus: ZNF81 (human) mapping to Xp11.23.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ZNF81 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF81 shRNA Plasmid (h): sc-77007-SH and ZNF81 shRNA (h) Lentiviral Particles: sc-77007-V as alternate gene silencing products.

For independent verification of ZNF81 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-77007A, sc-77007B and sc-77007C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF81 siRNA (h) is recommended for the inhibition of ZNF81 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF81 gene expression knockdown using RT-PCR Primer: ZNF81 (h)-PR: sc-77007-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.