

RBAK (N-14): sc-79007

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krueppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. RABK (RB-associated KRAB zinc finger), also known as ZNF769 (zinc finger protein 769), is a 714 amino acid protein that localizes to the nucleus and contains one KRAB domain and 16 C₂H₂-type zinc fingers. Expressed in liver, heart, kidney, placenta, pancreas, lung and bone, RBAK interacts with AR (androgen receptor) and Rb (retinoblastoma) and is thought to both promote AR-dependent transcription and repress E2F-dependent transcription.

REFERENCES

1. Skapek, S.X., Jansen, D., Wei, T.F., McDermott, T., Huang, W., Olson, E.N. and Lee, E.Y. 2000. Cloning and characterization of a novel Kruppel-associated box family transcriptional repressor that interacts with the retinoblastoma gene product, RB. *J. Biol. Chem.* 275: 7212-7223.
2. Gomes, I., Sharma, T.T., Edassery, S., Fulton, N., Mar, B.G. and Westbrook, C.A. 2002. Novel transcription factors in human CD34 antigen-positive hematopoietic cells. *Blood* 100: 107-119.
3. Hofman, K., Swinnen, J.V., Claessens, F., Verhoeven, G. and Heyns, W. 2003. The retinoblastoma protein-associated transcription repressor RBAK interacts with the androgen receptor and enhances its transcriptional activity. *J. Mol. Endocrinol.* 31: 583-596.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 608191. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Zhao, Y., Zhou, L., Liu, B., Deng, Y., Wang, Y., Wang, Y., Huang, W., Yuan, W., Wang, Z., Zhu, C., Liu, M., Wu, X. and Li, Y. 2006. ZNF325, a novel human zinc finger protein with a RBAK-like RB-binding domain, inhibits AP-1- and SRE-mediated transcriptional activity. *Biochem. Biophys. Res. Commun.* 346: 1191-1199.
6. So, A., Jeske, Y.W., Gordon, R.D., Duffy, D., Kelemen, L. and Stowasser, M. 2006. No evidence for coding region mutations in the retinoblastoma-associated Kruppel-associated box protein gene (RBAK) causing familial hyperaldosteronism type II. *Clin. Endocrinol.* 65: 829-831.
7. Jeske, Y.W., So, A., Kelemen, L., Sukor, N., Willys, C., Bulmer, B., Gordon, R.D., Duffy, D. and Stowasser, M. 2008. Examination of chromosome 7p22 candidate genes RBAK, PMS2 and GNA12 in familial hyperaldosteronism type II. *Clin. Exp. Pharmacol. Physiol.* 35: 380-385.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: RBAK (human) mapping to 7p22.1.

SOURCE

RBAK (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of RBAK of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79007 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

RBAK (N-14) is recommended for detection of RBAK of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RBAK siRNA (h): sc-76359, RBAK shRNA Plasmid (h): sc-76359-SH and RBAK shRNA (h) Lentiviral Particles: sc-76359-V.

Molecular Weight (predicted) of RBAK: 83 kDa.

Molecular Weight (observed) of RBAK: 89 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.