

ACADVL (C-15): sc-79198

BACKGROUND

ACADVL (acyl-Coenzyme A dehydrogenase, very long chain), also known as VLCAD, LCACD or ACAD6, is an inner mitochondrial membrane protein that belongs to the family of acyl-CoA dehydrogenases. Involved in lipid metabolism, ACADVL has catalytic activity toward esters of long chain and very long chain fatty acids and functions in the first step of the fatty acid β -oxidation pathway. Defects in the gene encoding ACADVL are the cause of very long chain acyl-CoA dehydrogenase deficiency (VLCAD deficiency), an autosomal recessive disease that leads to impaired long-chain fatty acid oxidation. VLCAD deficiency can give rise to three different phenotypes: a severe childhood form with a high incidence of mortality and cardiomyopathy; a mild childhood form with hypoketotic hypoglycemia and low mortality rates; and an adult form characterized by myoglobinuria and rhabdomyolysis. Two isoforms of ACADVL exist due to alternative splicing events.

REFERENCES

1. Mathur, A., Sims, H.F., Gopalakrishnan, D., Gibson, B., Rinaldo, P., Vockley, J., Hug, G. and Strauss, A.W. 1999. Molecular heterogeneity in very-long-chain acyl-CoA dehydrogenase deficiency causing pediatric cardiomyopathy and sudden death. *Circulation* 99: 1337-1343.
2. Andresen, B.S., Olpin, S., Poorthuis, B.J., Scholte, H.R., Vianey-Saban, C., Wanders, R., Ijlst, L., Morris, A., Pourfarzam, M., Bartlett, K., Baumgartner, E.R., deKlerk, J.B., Schroeder, L.D., Corydon, T.J., Lund, H., Winter, V., Bross, P., Bolund, L. and Gregersen, N. 1999. Clear correlation of genotype with disease phenotype in very-long-chain acyl-CoA dehydrogenase deficiency. *Am. J. Hum. Genet.* 64: 479-494.
3. Tong, M.K., Lam, C.S., Mak, T.W., Fu, M.Y., Ng, S.H., Wanders, R.J. and Tang, N.L. 2006. Very long-chain acyl-CoA dehydrogenase deficiency presenting as acute hypercapnic respiratory failure. *Eur. Respir. J.* 28: 447-450.
4. Gobin-Limballe, S., Djouadi, F., Aubey, F., Olpin, S., Andresen, B.S., Yamaguchi, S., Mandel, H., Fukao, T., Ruiter, J.P., Wanders, R.J., McAndrew, R., Kim, J.J. and Bastin, J. 2007. Genetic basis for correction of very-long-chain acyl-coenzyme A dehydrogenase deficiency by bezafibrate in patient fibroblasts: toward a genotype-based therapy. *Am. J. Hum. Genet.* 81: 1133-1143.
5. Spiekerkoetter, U. 2007. Effects of a fat load and exercise on asymptomatic VLCAD deficiency. *J. Inherit. Metab. Dis.* 30: 405.
6. Zia, A., Kolodny, E.H. and Pastores, G.M. 2007. Very long chain acyl-CoA dehydrogenase deficiency in a pair of mildly affected monozygotic twin sister in their late fifties. *J. Inherit. Metab. Dis.* 30: 817.
7. Goetzman, E.S., Wang, Y., He, M., Mohsen, A.W., Ninness, B.K. and Vockley, J. 2007. Expression and characterization of mutations in human very long-chain acyl-CoA dehydrogenase using a prokaryotic system. *Mol. Genet. Metab.* 91: 138-147.
8. Werdich, A.A., Baudenbacher, F., Dzhura, I., Jeyakumar, L.H., Kannankeril, P.J., Fleischer, S., LeGrone, A., Milatovic, D., Aschner, M., Strauss, A.W., Anderson, M.E. and Exil, V.J. 2007. Polymorphic ventricular tachycardia and abnormal Ca^{2+} handling in very-long-chain acyl-CoA dehydrogenase null mice. *Am. J. Physiol. Heart Circ. Physiol.* 292: H2202-H2211.

CHROMOSOMAL LOCATION

Genetic locus: ACADVL (human) mapping to 17p13.1; Acadvl (mouse) mapping to 11 B2-B5.

SOURCE

ACADVL (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of ACADVL of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79198 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ACADVL (C-15) is recommended for detection of ACADVL of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ACADVL siRNA (h): sc-72427, ACADVL siRNA (m): sc-72428, ACADVL shRNA Plasmid (h): sc-72427-SH, ACADVL shRNA Plasmid (m): sc-72428-SH, ACADVL shRNA (h) Lentiviral Particles: sc-72427-V and ACADVL shRNA (m) Lentiviral Particles: sc-72428-V.

Molecular Weight of ACADVL: 66 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.