

Fibulin-4 (E-19): sc-79240

BACKGROUND

Fibulin proteins contribute to normal development of elastic fiber systems in various types of organs that require elasticity, such as vasculature, lung and skin. Fibulin-4, also known as EFEMP2 (EGF-containing fibulin-like extracellular matrix protein 2), MBP1 or UPH1 is a 443 amino acid secreted protein that contains 6 EGF-like calcium-binding domains and belongs to the Fibulin family. Expressed ubiquitously with highest expression in heart, Fibulin-4 is essential for connective tissue development and elastic fiber formation, and may also play an important role in vascular patterning and collagen biosynthesis. Defects in the gene encoding Fibulin-4 are associated with autosomal recessive cutis laxa type I (CL type I), a connective tissue disorder that is inherited in both an autosomal dominant and an autosomal recessive manner and is characterized by inelastic tissue in all affected areas of the body.

REFERENCES

- Giltay, R., Timpl, R. and Kostka, G. 1999. Sequence, recombinant expression and tissue localization of two novel extracellular matrix proteins, fibulin-3 and fibulin-4. *Matrix Biol.* 18: 469-480.
- Katsanis, N., Venable, S., Smith, J.R. and Lupski, J.R. 2000. Isolation of a paralog of the Doyme honeycomb retinal dystrophy gene from the multiple retinopathy critical region on 11q13. *Hum. Genet.* 106: 66-72.
- Gallagher, W.M., Greene, L.M., Ryan, M.P., Sierra, V., Berger, A., Laurent-Puig, P. and Conseiller, E. 2001. Human fibulin-4: analysis of its biosynthetic processing and mRNA expression in normal and tumour tissues. *FEBS Lett.* 489: 59-66.
- Toto, L., Parodi, M.B., Baralle, F., Casari, G., Ravalico, G. and Romano, M. 2002. Genetic heterogeneity in Malattia Leventinese. *Clin. Genet.* 62: 399-403.
- Huchtagowder, V., Sausgruber, N., Kim, K.H., Angle, B., Marmorstein, L.Y. and Urban, Z. 2006. Fibulin-4: a novel gene for an autosomal recessive cutis laxa syndrome. *Am. J. Hum. Genet.* 78: 1075-1080.
- Xiang, Y., Sekine, T., Nakamura, H., Imajoh-Ohmi, S., Fukuda, H., Yudoh, K., Masuko-Hongo, K., Nishioka, K. and Kato, T. 2006. Fibulin-4 is a target of autoimmunity predominantly in patients with osteoarthritis. *J. Immunol.* 176: 3196-3204.
- McLaughlin, P.J., Chen, Q., Horiguchi, M., Starcher, B.C., Stanton, J.B., Broekelmann, T.J., Marmorstein, A.D., McKay, B., Mecham, R., Nakamura, T. and Marmorstein, L.Y. 2006. Targeted disruption of fibulin-4 abolishes elastogenesis and causes perinatal lethality in mice. *Mol. Cell. Biol.* 26: 1700-1709.

CHROMOSOMAL LOCATION

Genetic locus: EFEMP2 (human) mapping to 11q13.1; Efemp2 (mouse) mapping to 19 A.

SOURCE

Fibulin-4 (E-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Fibulin-4 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79240 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Fibulin-4 (E-19) is recommended for detection of Fibulin-4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Fibulin-4 (E-19) is also recommended for detection of Fibulin-4 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Fibulin-4 siRNA (h): sc-75017, Fibulin-4 siRNA (m): sc-75018, Fibulin-4 shRNA Plasmid (h): sc-75017-SH, Fibulin-4 shRNA Plasmid (m): sc-75018-SH, Fibulin-4 shRNA (h) Lentiviral Particles: sc-75017-V and Fibulin-4 shRNA (m) Lentiviral Particles: sc-75018-V.

Molecular Weight of Fibulin-4: 61 kDa.

Positive Controls: mouse skin extract: sc-364251.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **Fibulin-4 (2C8): sc-293492**, our highly recommended monoclonal alternative to Fibulin-4 (E-19).