

USP44 (F-15): sc-79329

BACKGROUND

The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. Through the use of a wide range of enzymes that can add or remove ubiquitin, the Ub pathway controls many intracellular processes such as signal transduction, transcriptional activation and cell cycle progression. USP44 (ubiquitin specific peptidase 44) is a 712 amino acid protein that contains one UBP-type zinc finger and belongs to the peptidase C19 family. Expressed in testis, USP44, catalyzes the conversion of a ubiquitin C-terminal thioester to a free ubiquitin and a thiol. USP44 is encoded by a gene that maps to human chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome.

CHROMOSOMAL LOCATION

Genetic locus: USP44 (human) mapping to 12q22; Usp44 (mouse) mapping to 10 C2.

SOURCE

USP44 (F-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of USP44 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79329 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

USP44 (F-15) is recommended for detection of USP44 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

USP44 (F-15) is also recommended for detection of USP44 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for USP44 siRNA (h): sc-76857, USP44 siRNA (m): sc-76858, USP44 shRNA Plasmid (h): sc-76857-SH, USP44 shRNA Plasmid (m): sc-76858-SH, USP44 shRNA (h) Lentiviral Particles: sc-76857-V and USP44 shRNA (m) Lentiviral Particles: sc-76858-V.


Molecular Weight of USP44: 81 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, Jurkat whole cell lysate: sc-2204 or K-562 whole cell lysate: sc-2203.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



USP44 (F-15): sc-79329. Western blot analysis of USP44 expression in Daudi (A), NIH/3T3 (B), Jurkat (C) and K-562 (D) whole cell lysates.



USP44 (F-15): sc-79329. Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing perinuclear and cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

- Zhang, Y., et al. 2012. USP44 regulates centrosome positioning to prevent aneuploidy and suppress tumorigenesis. *J. Clin. Invest.* 122: 4362-4374.
- Lu, H., et al. 2012. Transformation of human ovarian surface epithelial cells by Krüppel-like factor 8. *Oncogene* 33: 10-18.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **USP44 (G-2): sc-377203**, our highly recommended monoclonal alternative to USP44 (F-15).