

UTX (P-18): sc-79332

BACKGROUND

The Notch signaling pathway controls cellular interactions important for the specification of a variety of fates in both vertebrates and invertebrates. Key players in the Notch pathway are the TLE genes (for transducin-like enhancer of split, also designated ESG for enhancer of split groucho), which are human homologs of the *Drosophila* groucho gene. UTX (ubiquitously transcribed tetratricopeptide repeat, X chromosome) is a 1,401 amino acid nuclear protein that interacts with TLE1 (transducin-like enhancer of split 1) and, together, these proteins are thought to function as transcriptional repressors for a variety of targets. Expressed from a gene located on the inactive X chromosome, UTX functions as a histone demethylase that is involved in modulating the histone code (via demethylation of lysine residues on Histone H3) and in regulating Hox (homeobox) gene expression. UTX contains one JMJC domain and eight TPR repeats.

REFERENCES

- Greenfield, A., et al. 1998. The UTX gene escapes X inactivation in mice and humans. *Hum. Mol. Genet.* 7: 737-742.
- Grbavec, D., et al. 1999. Groucho/transducin-like enhancer of split (TLE) family members interact with the yeast transcriptional co-repressor SSN6 and mammalian SSN6-related proteins: implications for evolutionary conservation of transcription repression mechanisms. *Biochem. J.* 337: 13-17.
- Hong, S., et al. 2007. Identification of JmjC domain-containing UTX and JMJD3 as Histone H3 lysine 27 demethylases. *Proc. Natl. Acad. Sci. USA* 104: 18439-18444.
- Lan, F., et al. 2007. A Histone H3 lysine 27 demethylase regulates animal posterior development. *Nature* 449: 689-694.
- Agger, K., et al. 2007. UTX and JMJD3 are Histone H3K27 demethylases involved in HOX gene regulation and development. *Nature* 449: 731-734.
- Cho, Y.W., et al. 2007. PTIP associates with MLL3- and MLL4-containing Histone H3 lysine 4 methyltransferase complex. *J. Biol. Chem.* 282: 20395-20406.
- Issaeva, I., et al. 2007. Knockdown of ALR (MLL2) reveals ALR target genes and leads to alterations in cell adhesion and growth. *Mol. Cell. Biol.* 27: 1889-1903.
- Smith, E.R., et al. 2008. *Drosophila* UTX is a histone H3 Lys27 demethylase that colocalizes with the elongating form of RNA polymerase II. *Mol. Cell. Biol.* 28: 1041-1046.

CHROMOSOMAL LOCATION

Genetic locus: KDM6A (human) mapping to Xp11.3; Kdm6a (mouse) mapping to X A1.2.

SOURCE

UTX (P-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of UTX of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79332 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

UTX (P-18) is recommended for detection of UTX of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

UTX (P-18) is also recommended for detection of UTX in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for UTX siRNA (h): sc-76881, UTX siRNA (m): sc-76882, UTX shRNA Plasmid (h): sc-76881-SH, UTX shRNA Plasmid (m): sc-76882-SH, UTX shRNA (h) Lentiviral Particles: sc-76881-V and UTX shRNA (m) Lentiviral Particles: sc-76882-V.

Molecular Weight of UTX: 154 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **UTX (E-8): sc-514859** or **UTX (D-3): sc-514860**, our highly recommended monoclonal alternatives to UTX (P-18).