

xCT (N-19): sc-79359



The Power to Question

BACKGROUND

xCT, also known as SLC7A11 (solute carrier family 7, (cationic amino acid transporter, y⁺ system) member 11) or CCBR1, is a 501 amino acid multi-pass membrane protein that belongs to the polyamine-organocation superfamily of amino acid transporters. Existing as a disulfide-linked heterodimer with CD98, xCT functions as a member of a heteromeric Na⁺-independent anionic amino acid transport system that specifically facilitates the exchange of anionic amino acids for anionic forms of cystine and glutamate, thereby mediating the formation of glutathione within the cell. Due to its involvement in amino acid transport, xCT is associated with the pathogenesis of glioma-induced neurodegeneration and brain edema, as well as pancreatic cancer. The gene encoding xCT maps to human chromosome 4, which encodes nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes.

REFERENCES

1. Sato, H., et al. 1999. Cloning and expression of a plasma membrane cystine/glutamate exchange transporter composed of two distinct proteins. *J. Biol. Chem.* 274: 11455-11458.
2. Sato, H., et al. 2000. Molecular cloning and expression of human xCT, the light chain of amino acid transport system x_c⁻. *Antioxid. Redox Signal.* 2: 665-671.
3. Shih, A.Y. and Murphy, T.H. 2001. xCT cystine transporter expression in HEK293 cells: pharmacology and localization. *Biochem. Biophys. Res. Commun.* 282: 1132-1137.
4. Kim, J.Y., et al. 2001. Human cystine/glutamate transporter: cDNA cloning and upregulation by oxidative stress in glioma cells. *Biochim. Biophys. Acta* 1512: 335-344.
5. Bridges, C.C., et al. 2001. Structure, function, and regulation of human cystine/glutamate transporter in retinal pigment epithelial cells. *Invest. Ophthalmol. Vis. Sci.* 42: 47-54.
6. Bassi, M.T., et al. 2001. Identification and characterisation of human xCT that co-expresses, with 4F2 heavy chain, the amino acid transport activity system x_c⁻. *Pflugers Arch.* 442: 286-296.
7. Lo, M., et al. 2008. The x_c⁻ cystine/glutamate antiporter: a mediator of pancreatic cancer growth with a role in drug resistance. *Br. J. Cancer* 99: 464-472.
8. Savaskan, N.E., et al. 2008. Small interfering RNA-mediated xCT silencing in gliomas inhibits neurodegeneration and alleviates brain edema. *Nat. Med.* 14: 629-632.
9. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 607933. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: SLC7A11 (human) mapping to 4q28.3; Slc7a11 (mouse) mapping to 3 C.

SOURCE

xCT (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal cytoplasmic domain of xCT of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79359 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

xCT (N-19) is recommended for detection of xCT of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

xCT (N-19) is also recommended for detection of xCT in additional species, including equine, canine and porcine.

Suitable for use as control antibody for xCT siRNA (h): sc-76933, xCT siRNA (m): sc-76934, xCT shRNA Plasmid (h): sc-76933-SH, xCT shRNA Plasmid (m): sc-76934-SH, xCT shRNA (h) Lentiviral Particles: sc-76933-V and xCT shRNA (m) Lentiviral Particles: sc-76934-V.

Molecular Weight of xCT: 40 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.