

# ZNF24 (E-15): sc-79377

## BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF191 (zinc finger protein 191), also known as ZNF24, KOX17, ZSCAN3 or RSG-A, is a 368 amino acid nuclear protein that belongs to the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family. Expressed in tissues throughout the body with the exception of heart, ZNF191 functions as a transcriptional repressor for a variety of proteins, such as VEGF (vascular endothelial growth factor), and is thought to be important for early embryonic development and cell proliferation. ZNF191 contains four C<sub>2</sub>H<sub>2</sub>-type zinc fingers and one SCAN box domain and, upon DNA damage, may be phosphorylated by ATM or ATR.

## REFERENCES

1. Rousseau-Merck, M.F., et al. 1991. Chromosomal localization of two human zinc finger protein genes, ZNF24 (KOX17) and ZNF29 (KOX26), to 18q12 and 17p13-p12, respectively. *Genomics* 9: 154-161.
2. Shi, S.L., et al. 1998. Assignment of a novel zinc finger gene ZNF191 to human chromosome 18Q12.1 by human/rodent somatic cell hybrid panel and fluorescent *in situ* hybridization. *Shi Yan Sheng Wu Xue Bao* 31: 21-27.
3. Han, Z.G., et al. 1999. Molecular cloning of six novel Krüppel-like zinc finger genes from hematopoietic cells and identification of a novel *trans*-regulatory domain KRNB. *J. Biol. Chem.* 274: 35741-35748.
4. Williams, A.J., et al. 1999. The zinc finger-associated SCAN box is a conserved oligomerization domain. *Mol. Cell. Biol.* 19: 8526-8535.
5. Li, J.Z., et al. 2004. Establishment of transgenic mice carrying gene encoding human zinc finger protein 191. *World J. Gastroenterol.* 10: 264-267.
6. Li, J., et al. 2006. The zinc finger transcription factor 191 is required for early embryonic development and cell proliferation. *Exp. Cell Res.* 312: 3990-3998.
7. Harper, J., et al. 2007. Repression of vascular endothelial growth factor expression by the zinc finger transcription factor ZNF24. *Cancer Res.* 67: 8736-8741.

## CHROMOSOMAL LOCATION

Genetic locus: ZNF24 (human) mapping to 18q12.2; Zfp24 (mouse) mapping to 18 B1.

## SOURCE

ZNF24 (E-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ZNF24 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79377 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

ZNF24 (E-15) is recommended for detection of ZNF24 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ZNF24 (E-15) is also recommended for detection of ZNF24 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ZNF24 siRNA (h): sc-76969, ZNF24 siRNA (m): sc-76970, ZNF24 shRNA Plasmid (h): sc-76969-SH, ZNF24 shRNA Plasmid (m): sc-76970-SH, ZNF24 shRNA (h) Lentiviral Particles: sc-76969-V and ZNF24 shRNA (m) Lentiviral Particles: sc-76970-V.


Molecular Weight of ZNF24: 45 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, Jurkat nuclear extract: sc-2132 or THP-1 nuclear extract: sc-24963.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



ZNF191 (E-15): sc-79377. Western blot analysis of ZNF191 expression in Jurkat nuclear extract.



ZNF191 (E-15): sc-79377. Western blot analysis of ZNF191 expression in THP-1 nuclear extract.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.