

# ZNF569 (E-14): sc-79381



The Power to Question

## BACKGROUND

Zinc finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF569 (zinc finger protein 569), also known as ZAP1, is a 686 amino acid member of the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc finger protein family and is thought to be involved in transcriptional regulation. Localized to the nucleus, ZNF569 contains one KRAB domain and 18 C<sub>2</sub>H<sub>2</sub>-type zinc fingers through which it may convey DNA, RNA and protein binding capabilities.

## REFERENCES

1. Kato, N., Shimotohno, K., VanLeeuwen, D. and Cohen, M. 1990. Human proviral mRNAs down regulated in choriocarcinoma encode a zinc finger protein related to Krüppel. *Mol. Cell. Biol.* 10: 4401-4405.
2. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
3. Bray, P., Lichter, P., Thiesen, H.J., Ward, D.C. and Dawid, I.B. 1991. Characterization and mapping of human genes encoding zinc finger proteins. *Proc. Natl. Acad. Sci. USA* 88: 9563-9567.
4. Huebner, K., Druck, T., Croce, C.M. and Thiesen, H.J. 1991. Twenty-seven nonoverlapping zinc finger cDNAs from human T cells map to nine different chromosomes with apparent clustering. *Am. J. Hum. Genet.* 48: 726-740.
5. Lichter, P., Bray, P., Ried, T., Dawid, I.B. and Ward, D.C. 1992. Clustering of C<sub>2</sub>H<sub>2</sub> zinc finger motif sequences within telomeric and fragile site regions of human chromosomes. *Genomics* 13: 999-1007.
6. Englbrecht, C.C., Schoof, H. and Böhm, S. 2004. Conservation, diversification and expansion of C<sub>2</sub>H<sub>2</sub> zinc finger proteins in the *Arabidopsis thaliana* genome. *BMC Genomics* 5: 39-39.
7. Huang, X., Yuan, W., Huang, W., Bai, Y., Deng, Y., Zhu, C., Liang, P., Li, Y., Du, X., Liu, M., Wang, Y. and Wu, X. 2006. ZNF569, a novel KRAB-containing zinc finger protein, suppresses MAPK signaling pathway. *Biochem. Biophys. Res. Commun.* 346: 621-628.
8. Sjöblom, T., Jones, S., Wood, L.D., Parsons, D.W., Lin, J., Barber, T.D., Mandelker, D., Leary, R.J., Ptak, J., Silliman, N., Szabo, S., Buckhaults, P., Farrell, C., Meeh, P., Markowitz, S.D., Willis, J., Dawson, D., et al. 2006. The consensus coding sequences of human breast and colorectal cancers. *Science* 314: 268-274.

## CHROMOSOMAL LOCATION

Genetic locus: ZNF569 (human) mapping to 19q13.12.

## SOURCE

ZNF569 (E-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ZNF569 of human origin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79381 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

ZNF569 (E-14) is recommended for detection of ZNF569 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ZNF569 siRNA (h): sc-63253, ZNF569 shRNA Plasmid (h): sc-63253-SH and ZNF569 shRNA (h) Lentiviral Particles: sc-63253-V.


Molecular Weight of ZNF569: 80 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, HEK293 whole cell lysate: sc-45136 or A549 cell lysate: sc-2413.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



ZNF569 (E-14): sc-79381. Western blot analysis of ZNF569 expression in Hep G2 (A), HEK293 (B) and A549 (C) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.