

# CD100 (S-16): sc-79404

## BACKGROUND

Semaphorins comprise a family of phylogenetically conserved proteins containing a 500 amino acid region termed the "Sema domain" in the amino-terminus. The semaphorin family is subdivided into eight groups. CD100 (SEMA4D) is a transmembrane protein belonging to group IV of the semaphorin family. CD100 is expressed in both embryonic and adult kidney, heart, lung and neural tissues. Plexin-B1, a receptor expressed abundantly in fetal brain and kidney, binds CD100 with high affinity. CD100 is the only member of the semaphorin family that is expressed in the immune system. CD100 activates T cell proliferation and promotes B cell aggregation. CD100 regulates B cell response by negating an inhibitory signal from CD72 that normally diminishes B cell response. This signal is turned off when CD100 binds to CD72 and induces tyrosine dephosphorylation of CD72, followed by the dissociation of SHP-1 from CD72. The gene encoding human CD100 maps to chromosome 9q22.2.

## REFERENCES

1. Bougeret, C., et al. 1992. Increased surface expression of a newly identified 150 kDa dimer early after human T lymphocyte activation. *J. Immunol.* 148: 318-323.
2. Herold, C., et al. 1995. Activation signals are delivered through two distinct epitopes of CD100, a unique 150 kDa human lymphocyte surface structure previously defined by BB18 mAb. *Int. Immunol.* 7: 1-8.
3. Hall, K.T., et al. 1996. Human CD100, a novel leukocyte semaphorin that promotes B cell aggregation and differentiation. *Proc. Natl. Acad. Sci. USA* 93: 11780-11785.

## CHROMOSOMAL LOCATION

Genetic locus: SEMA4D (human) mapping to 9q22.2; Sema4d (mouse) mapping to 13 A5.

## SOURCE

CD100 (S-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of CD100 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79404 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## APPLICATIONS

CD100 (S-16) is recommended for detection of CD100 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CD100 (S-16) is also recommended for detection of CD100 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for CD100 siRNA (h): sc-42817, CD100 siRNA (m): sc-42818, CD100 shRNA Plasmid (h): sc-42817-SH, CD100 shRNA Plasmid (m): sc-42818-SH, CD100 shRNA (h) Lentiviral Particles: sc-42817-V and CD100 shRNA (m) Lentiviral Particles: sc-42818-V.


Molecular Weight of CD100: 150 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, SK-N-SH cell lysate: sc-2410 or WEHI-231 whole cell lysate: sc-2213.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



CD100 (S-16): sc-79404. Western blot analysis of CD100 expression in Jurkat whole cell lysate.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

**MONOS**  
Satisfaction  
Guaranteed

Try **CD100 (C-3): sc-390675** or **CD100 (D-5): sc-166367**, our highly recommended monoclonal alternatives to CD100 (S-16).