

PLAC4 (S-20): sc-79681

BACKGROUND

PLAC4 (placenta-specific 4), also known as D21S418E, is a 150 amino acid protein that is expressed in maternal plasma by placental syncytiotrophoblast and choriocarcinoma cells. PLAC4 is fetal derived, therefore it is specifically expressed during pregnancy and clears from maternal plasma within 24 hours after delivery. The PLAC4 gene is located on chromosome 21 and has four single nucleotide polymorphisms (SNPs) in its transcribed regions. This allows for PLAC4 to be a target gene in the noninvasive prenatal detection of trisomy 21 by allelic ratio analysis of placental expressed mRNA transcripts in maternal plasma. In a heterozygous fetus, both alleles of an SNP should be represented equally. An imbalance in allele frequency, giving a 2:1 ratio, suggests an additional copy of chromosome 21 and thus detection of trisomy 21. In addition, PLAC4 may be a useful target in the prenatal detection of Down syndrome.

REFERENCES

- Kido, S., Sakuragi, N., Bronner, M.P., Sayegh, R., Berger, R., Patterson, D. and Strauss J.F., 3rd. 1993. D21S418E identifies a cAMP-regulated gene located on chromosome 21q22.3 that is expressed in placental syncytiotrophoblast and choriocarcinoma cells. *Genomics* 17: 256-259.
- Lo, Y.M., Tsui, N.B., Chiu, R.W., Lau, T.K., Leung, T.N., Heung, M.M., Gerovassili, A., Jin, Y., Nicolaidis, K.H., Cantor, C.R. and Ding, C. 2007. Plasma placental RNA allelic ratio permits noninvasive prenatal chromosomal aneuploidy detection. *Nat. Med.* 13: 218-223.
- Lo, Y.M., Lun, F.M., Chan, K.C., Tsui, N.B., Chong, K.C., Lau, T.K., Leung, T.Y., Zee, B.C., Cantor, C.R. and Chiu, R.W. 2007. Digital PCR for the molecular detection of fetal chromosomal aneuploidy. *Proc. Natl. Acad. Sci. USA* 104: 13116-13121.
- Benachi, A. and Costa, J.M. 2007. Non-invasive prenatal diagnosis of fetal aneuploidies. *Lancet* 369: 440-442.
- Piriyapongsa, J., Polavarapu, N., Borodovsky, M. and McDonald, J. 2007. Exonization of the LTR transposable elements in human genome. *BMC Genomics* 8: 291.
- Jorgez, C.J., Dang, D.D., Wapner, R., Farina, A., Simpson, J.L. and Bischoff, F.Z. 2007. Elevated levels of total (maternal and fetal) β -globin DNA in maternal blood from first trimester pregnancies with trisomy 21. *Hum. Reprod.* 22: 2267-2272.
- Hahn, S., Zhong, X.Y. and Holzgreve, W. 2008. Recent progress in non-invasive prenatal diagnosis. *Semin. Fetal Neonatal. Med.* 13: 57-62.
- Go, A.T., Visser, A., Betsalel, O.T., van Vugt, J.M., Blankenstein, M.A. and Oudejans, C.B. 2008. Measurement of allelic-expression ratios in trisomy 21 placentas by quencher extension of heterozygous samples identified by partially denaturing HPLC. *Clin. Chem.* 54: 437-440.

CHROMOSOMAL LOCATION

Genetic locus: PLAC4 (human) mapping to 21q22.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

PLAC4 (S-20) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of PLAC4 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79681 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).


APPLICATIONS

PLAC4 (S-20) is recommended for detection of PLAC4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PLAC4 siRNA (h): sc-76166, PLAC4 shRNA Plasmid (h): sc-76166-SH and PLAC4 shRNA (h) Lentiviral Particles: sc-76166-V.

Molecular Weight of PLAC4: 17 kDa.

DATA



PLAC4 (S-20): sc-79681. Western blot analysis of PLAC4 expression in HEK293 (A), HeLa (B), JEG-3 (C), JAR (D), Jurkat (E) and A-431 (F) whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.