

sulfamidase (K-18): sc-79742

BACKGROUND

sulfatases are enzymes that hydrolyse a diverse range of sulfate esters. Deficiency of lysosomal sulfatases leads to human diseases characterized by the accumulation of either GAGs (glycosaminoglycans) or sulfolipids. sulfamidase, also known as HSS, SFMD, MPS3A or SGSH, is a 502 amino acid lysosome that belongs to the sulfatase family. It has been suggested that sulfamidase may be involved in the lysosomal degradation of heparan sulfate. Defects in the gene encoding sulfamidase are the cause of Sanfilippo syndrome A, an autosomal recessive lysosomal storage disease caused by impaired degradation of heparan sulfate. Sanfilippo syndrome A is characterized by severe central nervous system degeneration but relatively mild somatic manifestations.

REFERENCES

1. Scott, H.S., Blanch, L., Guo, X.H., Freeman, C., Orsborn, A., Baker, E., Sutherland, G.R., Morris, C.P. and Hopwood, J.J. 1995. Cloning of the sulphamidase gene and identification of mutations in Sanfilippo A syndrome. *Nat. Genet.* 11: 465-467.
2. Blanch, L., Weber, B., Guo, X.H., Scott, H.S. and Hopwood, J.J. 1997. Molecular defects in Sanfilippo syndrome type A. *Hum. Mol. Genet.* 6: 787-791.
3. Perkins, K.J., Byers, S., Yogalingam, G., Weber, B. and Hopwood, J.J. 1999. Expression and characterization of wildtype and mutant recombinant human sulfamidase. Implications for Sanfilippo (Mucopolysaccharidosis IIIA) syndrome. *J. Biol. Chem.* 274: 37193-37199.
4. Muschol, N., Storch, S., Ballhausen, D., Beesley, C., Westermann, J.C., Gal, A., Ullrich, K., Hopwood, J.J., Winchester, B. and Braulke, T. 2004. Transport, enzymatic activity, and stability of mutant sulfamidase (SGSH) identified in patients with mucopolysaccharidosis type III A. *Hum. Mutat.* 23: 559-566.
5. Mason, K.E., Meikle, P.J., Hopwood, J.J. and Fuller, M. 2006. Characterization of sulfated oligosaccharides in mucopolysaccharidosis type IIIA by electrospray ionization mass spectrometry. *Anal. Chem.* 78: 4534-4542.
6. Di Natale, P., Pontarelli, G., Villani, G.R. and Di Domenico, C. 2006. Gene symbol: SGSH. Disease: Sanfilippo type A syndrome, mucopolysaccharidosis IIIA. *Hum. Genet.* 119: 679.
7. Lau, A.A., Hemsley, K.M., Meedeniya, A., Robinson, A.J. and Hopwood, J.J. 2006. Directed differentiation and characterization of genetically modified embryonic stem cells for therapy. *Methods Mol. Biol.* 329: 471-484.
8. Fraldi, A., Biffi, A., Lombardi, A., Visigalli, I., Pepe, S., Settembre, C., Nusco, E., Auricchio, A., Naldini, L., Ballabio, A. and Cosma, M.P. 2007. SUMF1 enhances sulfatase activities *in vivo* in five sulfatase deficiencies. *Biochem. J.* 403: 305-312.
9. Urayama, A., Grubb, J.H., Sly, W.S. and Banks, W.A. 2008. Mannose 6-phosphate receptor-mediated transport of sulfamidase across the blood-brain barrier in the newborn mouse. *Mol. Ther.* 16: 1261-1266.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

CHROMOSOMAL LOCATION

Genetic locus: SGSH (human) mapping to 17q25.3; Sgsh (mouse) mapping to 11 E2.

SOURCE

sulfamidase (K-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of sulfamidase of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79742 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

sulfamidase (K-18) is recommended for detection of sulfamidase of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

sulfamidase (K-18) is also recommended for detection of sulfamidase in additional species, including equine, canine and bovine.

Suitable for use as control antibody for sulfamidase siRNA (h): sc-76607, sulfamidase siRNA (m): sc-76608, sulfamidase shRNA Plasmid (h): sc-76607-SH, sulfamidase shRNA Plasmid (m): sc-76608-SH, sulfamidase shRNA (h) Lentiviral Particles: sc-76607-V and sulfamidase shRNA (m) Lentiviral Particles: sc-76608-V.

Molecular Weight of sulfamidase: 62 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.