

# TRIM11 (P-17): sc-79759

## BACKGROUND

The tripartite motif (TRIM) family of proteins are characterized by a conserved TRIM domain that includes a coiled-coil region, a B box-type zinc finger, one RING finger and three zinc-binding domains. TRIM11 (tripartite motif-containing 11), also known as BIA1 or RNF92, is a 468 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one RING-type zinc finger, one SPRY domain and one B box-type zinc finger. Expressed ubiquitously, TRIM11 is thought to function as an E3 ubiquitin ligase that may regulate the intracellular level of select proteins via control of the proteasomal degradation pathway. Multiple isoforms of TRIM11 exist due to alternative splicing events.

## REFERENCES

1. Reymond, A., Meroni, G., Fantozzi, A., Merla, G., Cairo, S., Luzi, L., Riganelli, D., Zanaria, E., Messali, S., Cainarca, S., Guffanti, A., Minucci, S., Pelicci, P.G. and Ballabio, A. 2001. The tripartite motif family identifies cell compartments. *EMBO J.* 20: 2140-2151.
2. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 607868. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Meroni, G. and Diez-Roux, G. 2005. TRIM/RBCC, a novel class of "single protein RING finger" E3 ubiquitin ligases. *Bioessays* 27: 1147-1157.
4. Ishikawa, H., Tachikawa, H., Miura, Y. and Takahashi, N. 2006. TRIM11 binds to and destabilizes a key component of the activator-mediated cofactor complex (ARC105) through the ubiquitin-proteasome system. *FEBS Lett.* 580: 4784-4792.
5. Sardiello, M., Cairo, S., Fontanella, B., Ballabio, A. and Meroni, G. 2008. Genomic analysis of the TRIM family reveals two groups of genes with distinct evolutionary properties. *BMC Evol. Biol.* 8: 225.
6. Hong, S.J., Chae, H., Lardaro, T., Hong, S. and Kim, K.S. 2008. TRIM11 increases expression of dopamine  $\beta$ -hydroxylase gene by interacting with Phox2b. *Biochem. Biophys. Res. Commun.* 368: 650-655.
7. Bowie, A.G. 2008. TRIM-ing down Tolls. *Nat. Immunol.* 9: 348-350.
8. Ozato, K., Shin, D.M., Chang, T.H. and Morse, H.C. 2008. TRIM family proteins and their emerging roles in innate immunity. *Nat. Rev. Immunol.* 8: 849-860.

## CHROMOSOMAL LOCATION

Genetic locus: TRIM11 (human) mapping to 1q42.13; Trim11 (mouse) mapping to 11 B1.3.

## SOURCE

TRIM11 (P-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TRIM11 of human origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79759 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

TRIM11 (P-17) is recommended for detection of TRIM11 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TRIM11 (P-17) is also recommended for detection of TRIM11 in additional species, including canine and bovine.

Suitable for use as control antibody for TRIM11 siRNA (h): sc-76734, TRIM11 siRNA (m): sc-76735, TRIM11 shRNA Plasmid (h): sc-76734-SH, TRIM11 shRNA Plasmid (m): sc-76735-SH, TRIM11 shRNA (h) Lentiviral Particles: sc-76734-V and TRIM11 shRNA (m) Lentiviral Particles: sc-76735-V.

Molecular Weight of TRIM11: 53 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.