

p-Akt1/2/3 (Ser 473)-R: sc-7985-R

BACKGROUND

The serine/threonine kinase Akt family contains several members, including Akt1 (also designated PKB or RacPK), Akt2 (also designated PKB β or RacPK- β) and Akt 3 (also designated PKB γ or thymoma viral proto-oncogene 3), which exhibit sequence homology with the protein kinase A and C families and are encoded by the c-Akt proto-oncogene. All members of the Akt family have a Pleckstrin homology domain. Akt1 and Akt2 are activated by PDGF stimulation. This activation is dependent on PDGFR- β tyrosine residues 740 and 751, which bind the subunit of the phosphatidylinositol 3-kinase (PI 3-kinase) complex. Activation of Akt1 by Insulin or Insulin-growth factor-1(IGF-1) results in phosphorylation of both Thr 308 and Ser 473. Akt proteins become phosphorylated and activated in Insulin/IGF-1-stimulated cells by an upstream kinase(s), and the activation of Akt1 and Akt2 is inhibited by the PI kinase inhibitor Wortmannin. Taken together, this data strongly suggests that the protein signals downstream of the PI kinases. Akt3 is phosphorylated on a serine residue in response to Insulin. However, the activation of Akt3 by Insulin is inhibited by prior activation of protein kinase C via a mechanism that does not require the presence of the PH domain. Akt3 is expressed in 3T3-L1 fibroblasts, adipocytes and skeletal muscle and may be involved in various biological processes, including adipocyte and muscle differentiation, glycogen synthesis, glucose uptake, apoptosis and cellular proliferation.

SOURCE

p-Akt1/2/3 (Ser 473)-R is a rabbit polyclonal antibody raised against a short amino acid sequence containing Ser 473 phosphorylated Akt1 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-7985 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

p-Akt1/2/3 (Ser 473)-R is recommended for detection of Ser 473 phosphorylated Akt1 and correspondingly Ser 474 phosphorylated Akt2 and correspondingly Ser 472 phosphorylated Akt3 of mouse, rat, human and *Xenopus laevis* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

p-Akt1/2/3 (Ser 473)-R is also recommended for detection of correspondingly phosphorylated Akt1, Akt2 and Akt3 in additional species, including bovine, porcine and avian.

Molecular Weight of p-Akt1: 62 kDa.

Molecular Weight of p-Akt2: 56 kDa.


Molecular Weight of p-Akt3: 62 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201, Jurkat whole cell lysate: sc-2204 or HeLa + heat shock cell lysate: sc-2272.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



Western blot analysis of Akt1/2/3 phosphorylation in untreated (A, D), calyculin treated (B, E) and calyculin and lambda protein phosphatase (sc-200312A) treated (C, F) Jurkat whole cell lysates. Antibodies tested include p-Akt1/2/3 (Ser 473)-R: sc-7985-R (A, B, C) and Akt1 (C-20): sc-1618 (D, E, F).



p-Akt1/2/3 (Ser 473)-R: sc-7985-R. Immunofluorescence staining of methanol-fixed A-431 cells showing nuclear localization.

SELECT PRODUCT CITATIONS

- Contos, J.J., et al. 2002. Characterization of Ipa₂ (EDG-4) and Ipa₁/Ipa₂ (EDG-2/EDG-4) lysophosphatidic acid receptor knockout mice: signaling deficits without obvious phenotypic abnormality attributable to Ipa₂. *Mol. Cell. Biol.* 22: 6921-6929.
- Zhang, J., et al. 2015. Promotion of dental pulp cell migration and pulp repair by a bioceramic putty involving FGFR-mediated signaling pathways. *J. Dent. Res.* 94: 853-862.
- Pereira, J.K., et al. 2015. Molecular effects of the phosphatidylinositol-3-kinase inhibitor NVP-BKM120 on T and B-cell acute lymphoblastic leukaemia. *Eur. J. Cancer* 51: 2076-2085.
- Das, A., et al. 2015. A novel component from citrus, ginger, and mushroom family exhibits antitumor activity on human meningioma cells through suppressing the Wnt/ β -catenin signaling pathway. *Tumour Biol.* 36: 7027-7034.
- Das, A., et al. 2015. Synergistic effects of crizotinib and temozolomide in experimental FIG-ROS1 fusion-positive glioblastoma. *Cancer Growth Metastasis* 8: 51-60.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
Satisfation
Guaranteed

Try **p-Akt1/2/3 (C-11): sc-514032** or **p-Akt1/2/3 (11E6): sc-81433**, our highly recommended monoclonal alternatives to p-Akt1/2/3 (Ser 473).