

Cdc2L5 (S-19): sc-79895

BACKGROUND

The eukaryotic cell division cycle consists of a number of gene-controlled sequences that involve cyclin-dependent kinases (Cdks) and cell division control (Cdc) proteins. Cdc2L5 (cell division cycle 2-like 5), also known as CHED or CDC2L, is a member of the cyclin-dependent serine/threonine protein kinase family that plays a role in cell differentiation and apoptosis. Expressed in liver, brain and muscle, Cdc2L5 is thought to control key events in the mitotic cycle and may be involved in the development of blood cells. Due to its role in mitosis, overexpression of Cdc2L5 is implicated in the development of neuroblastoma and glioblastoma tumors, suggesting a possible role for Cdc2L5 in carcinogenesis. Cdc2L5 contains one protein kinase domain and is expressed as two different isoforms produced by alternative splicing events.

REFERENCES

1. Lapidot-Lifson, Y., Patinkin, D., Prody, C.A., Ehrlich, G., Seidman, S., Ben-Aziz, R., Benseler, F., Eckstein, F., Zakut, H. and Soreq, H. 1992. Cloning and antisense oligodeoxynucleotide inhibition of a human homolog of Cdc2 required in hematopoiesis. *Proc. Natl. Acad. Sci. USA* 89: 579-583.
2. Marqués, F., Moreau, J.L., Peaucellier, G., Lozano, J.C., Schatt, P., Picard, A., Callebaut, I., Perret, E. and Genevière, A.M. 2000. A new subfamily of high molecular mass Cdc2-related kinases with PITAI/VRE motifs. *Biochem. Biophys. Res. Commun.* 279: 832-837.
3. Kristich, C.J. and Ordal, G.W. 2002. *Bacillus subtilis* CheD is a chemoreceptor modification enzyme required for chemotaxis. *J. Biol. Chem.* 277: 25356-25362.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603309. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Even, Y., Durieux, S., Escande, M.L., Lozano, J.C., Peaucellier, G., Weil, D. and Genevière, A.M. 2006. Cdc2L5, a Cdk-like kinase with RS domain, interacts with the ASF/SF2-associated protein p32 and affects splicing *in vivo*. *J. Cell. Biochem.* 99: 890-904.
6. Chen, H.H., Wong, Y.H., Geneviere, A.M. and Fann, M.J. 2007. Cdk13/Cdc2L5 interacts with L-type cyclins and regulates alternative splicing. *Biochem. Biophys. Res. Commun.* 354: 735-740.

CHROMOSOMAL LOCATION

Genetic locus: CDK13 (human) mapping to 7p14.1; Cdk13 (mouse) mapping to 13 A2.

SOURCE

CDC2L5 (S-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CDC2L5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79895 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Cdc2L5 (S-19) is recommended for detection of Cdc2L5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Cdks family members.

Cdc2L5 (S-19) is also recommended for detection of Cdc2L5 in additional species, including bovine.

Suitable for use as control antibody for Cdc2L5 siRNA (h): sc-72835, Cdc2L5 siRNA (m): sc-72836, Cdc2L5 shRNA Plasmid (h): sc-72835-SH, Cdc2L5 shRNA Plasmid (m): sc-72836-SH, Cdc2L5 shRNA (h) Lentiviral Particles: sc-72835-V and Cdc2L5 shRNA (m) Lentiviral Particles: sc-72836-V.

Molecular Weight of Cdc2L5: 165 kDa.

Positive Controls: HeLa nuclear extract: sc-2120.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Cdc2L5 (ZZ08): sc-81837**, our highly recommended monoclonal alternative to Cdc2L5 (S-19).