

COX5b (N-17): sc-79919

BACKGROUND

The cytochrome c oxidase (COX) family of proteins function as the final electron donor in the respiratory chain to drive a proton gradient across the inner mitochondrial membrane, ultimately resulting in the production of water. The mammalian COX apoenzyme is a dimer, with each monomer consisting of 13 subunits, some of which are mitochondrial and some of which are nuclear. Found in the inner mitochondrial membrane, COX5 is the heme A-containing chain of the oxidase family that converts one molecule of oxygen and four molecules of hydrogen to two molecules of water. Two isoforms of COX5 exist, COX5a and COX5b. When oxygen levels within the cell are high, transcription of COX5a (the aerobic isoform) is upregulated as the rate of cellular respiration increases. Conversely, when oxygen levels are low, COX5b (the hypoxic isoform) transcription increases and functions to maximize the turnover rate of the COX apoenzyme.

REFERENCES

- Hodge, M.R., et al. 1989. Inverse regulation of the yeast COX5 genes by oxygen and heme. *Mol. Cell. Biol.* 9: 1958-1964.
- Allen, L.A., et al. 1995. Isoforms of yeast cytochrome c oxidase subunit V affect the binuclear reaction center and alter the kinetics of interaction with the isoforms of yeast cytochrome c. *J. Biol. Chem.* 270: 110-118.
- Bachman, N.J., et al. 1996. Phylogenetic footprinting of the human cytochrome c oxidase subunit VB promoter. *Arch. Biochem. Biophys.* 333: 152-162.
- Burke, P.V. and Poyton, R.O. 1998. Structure/function of oxygen-regulated isoforms in cytochrome c oxidase. *J. Exp. Biol.* 201: 1163-1175.
- David, P.S. and Poyton, R.O. 2005. Effects of a transition from normoxia to anoxia on yeast cytochrome c oxidase and the mitochondrial respiratory chain: implications for hypoxic gene induction. *Biochim. Biophys. Acta* 1709: 169-180.
- Dang, C., et al. 2006. Identification of dysregulated genes in cutaneous squamous cell carcinoma. *Oncol. Rep.* 16: 513-519.
- Yamasaki, H., et al. 2007. Regulation of copper homeostasis by micro-RNA in *Arabidopsis*. *J. Biol. Chem.* 282: 16369-16378.
- Guescini, M., et al. 2007. Fine needle aspiration coupled with real-time PCR: a painless methodology to study adaptive functional changes in skeletal muscle. *Nutr. Metab. Cardiovasc. Dis.* 17: 383-393.
- Fukuda, R., et al. 2007. HIF-1 regulates cytochrome oxidase subunits to optimize efficiency of respiration in hypoxic cells. *Cell* 129: 111-122.

CHROMOSOMAL LOCATION

Genetic locus: COX5B (human) mapping to 2q11.2; Cox5b (mouse) mapping to 1 B.

SOURCE

COX5b (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of COX5b of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79919 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

COX5b (N-17) is recommended for detection of COX5b of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for COX5b siRNA (h): sc-72982, COX5b siRNA (m): sc-72983, COX5b shRNA Plasmid (h): sc-72982-SH, COX5b shRNA Plasmid (m): sc-72983-SH, COX5b shRNA (h) Lentiviral Particles: sc-72982-V and COX5b shRNA (m) Lentiviral Particles: sc-72983-V.

Molecular Weight of COX5b: 14 kDa.

Positive Controls: HeLa nuclear extract: sc-2120.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **COX5b (C-5): sc-374416** or **COX5b (B-2): sc-374417**, our highly recommended monoclonal alternatives to COX5b (N-17).