

EF-Ts (L-20): sc-79934

BACKGROUND

EF-Ts, also known as TSFM (Ts translation elongation factor, mitochondrial) or COXPD3, is a 325 amino acid protein that is one of 13 mitochondrial-encoded proteins that work together during the elongation phase of protein biosynthesis on the ribosome. Expressed ubiquitously with highest levels present in liver, kidney and skeletal muscle, EF-Ts associates with EF-Tu, a multidomain GTPase with essential functions in translation, and, via this interaction, facilitates the exchange of GDP for GTP, thereby inducing protein elongation. Mutations in the gene encoding EF-Ts are the cause of combined oxidative phosphorylation deficiency type 3 (COXPD3), a condition characterized by defects in the mitochondrial oxidative phosphorylation system and often characterized by severe metabolic acidosis with encephalomyopathy or with hypertrophic cardiomyopathy. Multiple isoforms of EF-Ts exist due to alternative splicing events.

REFERENCES

- Xin, H., Worjiax, V., Burkhart, W. and Spremulli, L.L. 1995. Cloning and expression of mitochondrial translational elongation factor Ts from bovine and human liver. *J. Biol. Chem.* 270: 17243-17249.
- Benkowski, L.A., Takemoto, C., Ott, G., Beikman, M., Ueda, T., Watanabe, K., Sprinzl, M. and Spremulli, L.L. 1995. Interaction of mitochondrial elongation factors Tu.Ts with aminoacyl-tRNA. *Nucleic Acids Symp. Ser.* 33: 163-166.
- Vernon, J.L., Burr, P.C., Wiley, J.E. and Farwell, M.A. 2000. Assignment of the mitochondrial translation elongation factor Ts gene (TSFM) to human chromosome 12 bands q13→q14 by *in situ* hybridization and with somatic cell hybrids. *Cytogenet. Cell Genet.* 89: 145-146.
- Smeitink, J.A., Elpeleg, O., Antonicka, H., Diepstra, H., Saada, A., Smits, P., Sasarman, F., Vriend, G., Jacob-Hirsch, J., Shaag, A., Rechavi, G., Welling, B., Horst, J., Rodenburg, R.J., van den Heuvel, B. and Shoubridge, E.A. 2006. Distinct clinical phenotypes associated with a mutation in the mitochondrial translation elongation factor EFTs. *Am. J. Hum. Genet.* 79: 869-877.
- Antonicka, H., Sasarman, F., Kennaway, N.G. and Shoubridge, E.A. 2006. The molecular basis for tissue specificity of the oxidative phosphorylation deficiencies in patients with mutations in the mitochondrial translation factor EFG1. *Hum. Mol. Genet.* 15: 1835-1846.
- Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 604723. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: TSFM (human) mapping to 12q14.1; Tsfm (mouse) mapping to 10 D3.

SOURCE

EF-Ts (L-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of EF-Ts of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-79934 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

EF-Ts (L-20) is recommended for detection of EF-Ts of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

EF-Ts (L-20) is also recommended for detection of EF-Ts in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for EF-Ts siRNA (h): sc-77243, EF-Ts siRNA (m): sc-77244, EF-Ts shRNA Plasmid (h): sc-77243-SH, EF-Ts shRNA Plasmid (m): sc-77244-SH, EF-Ts shRNA (h) Lentiviral Particles: sc-77243-V and EF-Ts shRNA (m) Lentiviral Particles: sc-77244-V.

Molecular Weight of EF-Ts: 34 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.