

PKC (MC5): sc-80

BACKGROUND

Members of the protein kinase C (PKC) family play a key regulatory role in a variety of cellular functions, including cell growth and differentiation, gene expression, hormone secretion and membrane function. PKCs were originally identified as serine/threonine protein kinases whose activity was dependent on calcium and phospholipids. Diacylglycerols (DAG) and tumor promoting phorbol esters bind to and activate PKC. PKCs can be subdivided into at least two major classes, including conventional (c) PKC isoforms (α , β I, β II and γ) and novel (n) PKC isoforms (δ , ϵ , ζ , η , θ , λ , ι , μ and ν). Patterns of expression for each PKC isoform differ among tissues and PKC family members exhibit clear differences in their cofactor dependencies. For instance, the kinase activities of PKC δ and ϵ are independent of Ca^{2+} . On the other hand, most of the other PKC members possess phorbol ester-binding activities and kinase activities.

SOURCE

PKC (MC5) is a mouse monoclonal antibody epitope mapping within the hinge region (residues 292-317) of protein kinase C.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

PKC (MC5) is available conjugated to agarose (sc-80 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-80 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-80 PE), fluorescein (sc-80 FITC), Alexa Fluor[®] 488 (sc-80 AF488), Alexa Fluor[®] 546 (sc-80 AF546), Alexa Fluor[®] 594 (sc-80 AF594) or Alexa Fluor[®] 647 (sc-80 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-80 AF680) or Alexa Fluor[®] 790 (sc-80 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

PKC (MC5) is recommended for detection of PKC α , PKC β and PKC γ of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

PKC (MC5) is also recommended for detection of PKC α , PKC β and PKC γ in additional species, including bovine.

Suitable for use as control antibody for PKC siRNA (h): sc-29449, PKC shRNA Plasmid (h): sc-29449-SH and PKC shRNA (h) Lentiviral Particles: sc-29449-V.

Molecular Weight of PKC: 80 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, SUP-T1 whole cell lysate: sc-364796 or C6 whole cell lysate: sc-364373.


RESEARCH USE

For research use only, not for use in diagnostic procedures.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



PKC (MC5): sc-80. Western blot analysis of PKC expression in Jurkat (A), SUP-T1 (B), Caki-1 (C) and C6 (D) whole cell lysates.



PKC (MC5): sc-80. Immunoperoxidase staining of formalin fixed, paraffin-embedded human gall bladder tissue showing membrane and cytoplasmic staining of glandular cells at low (A) and high (B) magnification. Kindly provided by The Swedish Human Protein Atlas (HPA) program.

SELECT PRODUCT CITATIONS

- Kawakami, Y., et al. 1994. Tyrosine phosphorylation and activation of Bruton tyrosine kinase upon Fc ϵ RI cross-linking. *Mol. Cell. Biol.* 14: 5108-5113.
- Pilotto, S., et al. 2016. LSD1/KDM1A mutations associated to a newly described form of intellectual disability impair demethylase activity and binding to transcription factors. *Hum. Mol. Genet.* 25: 2578-2587.
- Li, L.H., et al. 2017. A synthetic cationic antimicrobial peptide inhibits inflammatory response and the NLRP3 inflammasome by neutralizing LPS and ATP. *PLoS ONE* 12: e0182057.
- Nguyen Dinh Cat, A., et al. 2018. Vascular dysfunction in obese diabetic db/db mice involves the interplay between aldosterone/mineralocorticoid receptor and Rho kinase signaling. *Sci. Rep.* 8: 2952.
- Mateus Gonçalves, L., et al. 2019. Amino acid restriction increases β -cell death under challenging conditions. *J. Cell. Physiol.* 234: 16679-16684.
- Ruiz-Pastor, M.J., et al. 2021. Decrease in DHA and other fatty acids correlates with photoreceptor degeneration in retinitis pigmentosa. *Exp. Eye Res.* 209: 108667.
- Chen, X., et al. 2022. Stearic acid induces CCK and GLP-1 upregulation via GPR120/PLC- β , leading to reduced appetite in Hu sheep fed with rice straw. *Front. Vet. Sci.* 9: 948074.
- Clemons, M.R., et al. 2023. The rod synapse in aging wildtype and Dscaml1 mutant mice. *PLoS ONE* 18: e0290257.
- Nadel, G., et al. 2024. Phosphorylation of PP2Ac by PKC is a key regulatory step in the PP2A-switch-dependent AKT dephosphorylation that leads to apoptosis. *Cell Commun. Signal.* 22: 154.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.