

# Bystin (clone 19): sc-80001

## BACKGROUND

Bystin (or Bystin-like) is an evolutionarily conserved protein from yeast to humans. It localizes to the nucleolus and cytoplasm. Nuclear Bystin associates with the 40S ribosomal subunit and cytoplasmic Bystin directly binds Trophinin and Tastin, facilitating the adhesion of cells expressing these proteins. Knock-down of Bystin inhibits cell proliferation and delays RNA processing of the 18S component of the 40S ribosomal subunit. This suggests that Bystin plays an important role in ribosome biogenesis. In addition, embryo development is arrested around the 16-cell stage when Bystin expression is knocked down. This implies that Bystin is important for the development of preimplantation embryos. Bystin is also overexpressed in human cancers providing further evidence suggesting that it participates in cell proliferation.

## REFERENCES

1. Fukuda, M.N. and Nozawa, S. 2000. Trophinin, Tastin, and Bystin: a complex mediating unique attachment between trophoblastic and endometrial epithelial cells at their respective apical cell membranes. *Semin. Reprod. Endocrinol.* 17: 229-234.
2. Aoki, R. and Fukuda, M.N. 2001. Recent molecular approaches to elucidate the mechanism of embryo implantation: Trophinin, Bystin, and Tastin as molecules involved in the initial attachment of blastocysts to the uterus in humans. *Semin. Reprod. Med.* 18: 265-271.
3. Sheng, J., Yang, S., Xu, L., Wu, C., Wu, X., Li, A., Yu, Y., Ni, H., Fukuda, M. and Zhou, J. 2004. Bystin as a novel marker for reactive astrocytes in the adult rat brain following injury. *Eur. J. Neurosci.* 20: 873-884.
4. Aoki, R., Suzuki, N., Paria, B.C., Sugihara, K., Akama, T.O., Raab, G., Miyoshi, M., Nadano, D. and Fukuda, M.N. 2006. The Bysl gene product, Bystin, is essential for survival of mouse embryos. *FEBS Lett.* 580: 6062-6068.
5. Ma, L., Yin, M., Wu, X., Wu, C., Yang, S., Sheng, J., Ni, H., Fukuda, M.N. and Zhou, J. 2006. Expression of Trophinin and Bystin identifies distinct cell types in the germinal zones of adult rat brain. *Eur. J. Neurosci.* 23: 2265-2276.
6. Ayala, G.E., Dai, H., Li, R., Ittmann, M., Thompson, T.C., Rowley, D. and Wheeler, T.M. 2006. Bystin in perineural invasion of prostate cancer. *Prostate* 66: 266-272.
7. Miyoshi, M., Okajima, T., Matsuda, T., Fukuda, M.N. and Nadano, D. 2007. Bystin in human cancer cells: intracellular localization and function in ribosome biogenesis. *Biochem. J.* 404: 373-381.
8. Adachi, K., Soeta-Saneyoshi, C., Sagara, H. and Iwakura, Y. 2007. Crucial role of BYSL in mammalian preimplantation development as an integral factor for 40S ribosome biogenesis. *Mol. Cell. Biol.* 27: 2202-2214.
9. Fukuda, M.N. and Sugihara, K. 2007. Signal transduction in human embryo implantation. *Cell Cycle* 6: 1153-1156.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## CHROMOSOMAL LOCATION

Genetic locus: BYSL (human) mapping to 6p21.1.

## SOURCE

Bystin (clone 19) is a mouse monoclonal antibody raised against a peptide corresponding to amino acids 215-223 of Bystin of human origin.

## PRODUCT

Each vial contains 200 µg IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

Bystin (clone 19) is recommended for detection of Bystin of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).


Suitable for use as control antibody for Bystin siRNA (h): sc-62030, Bystin shRNA Plasmid (h): sc-62030-SH and Bystin shRNA (h) Lentiviral Particles: sc-62030-V.

Molecular Weight of Bystin: 50 kDa.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 3) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



Bystin (clone 19): sc-80001. Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing cytoplasmic staining.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.