

MAP-1B (AA9): sc-80014

BACKGROUND

Microtubules, the primary component of the cytoskeletal network, interact with proteins called microtubule-associated proteins (MAPs). The microtubule-associated proteins can be divided into two groups, structural and dynamic. The structural microtubule-associated proteins, MAP-1A, MAP-1B, MAP-2A, MAP-2B and MAP-2C, stimulate Tubulin assembly, enhance microtubule stability and influence the spatial distribution of microtubules within cells. Both MAP-1 and, to a greater extent, MAP-2 have been implicated as agents of microtubule depolymerization by suppressing the dynamic instability of the microtubules. The suppression of microtubule dynamic instability by the MAP proteins is thought to be associated with phosphorylation of the MAPs.

REFERENCES

1. Sloboda, R.D., Dentler, W.L. and Rosenbaum, J.L. 1976. Microtubule-associated proteins and the stimulation of tubulin assembly *in vitro*. *Biochemistry* 15: 4497-4505.
2. Murphy, D.B., Johnson, K.A. and Borisy, G.G. 1977. Role of Tubulin-associated proteins in microtubule nucleation and elongation. *J. Mol. Biol.* 117: 33-52.
3. Hasegawa, M., Arai, T. and Ihara, Y. 1990. Immunochemical evidence that fragments of phosphorylated MAP-5 (MAP-1B) are bound to neurofibrillary tangles in Alzheimer's disease. *Neuron* 4: 909-918.
4. MacRae, T.H. 1992. Towards an understanding of microtubule function and cell organization: an overview. *Biochem. Cell Biol.* 70: 835-841.
5. Davis, R.J. 1993. The mitogen-activated protein kinase signal transduction pathway. *J. Biol. Chem.* 268: 14553-14556.
6. Maccioni, R.B. and Cambiasso, V. 1995. Role of microtubule-associated proteins in the control of microtubule assembly. *Physiol. Rev.* 75: 835-864.
7. Dhamodharan, R. and Wadsworth, P. 1995. Modulation of microtubule dynamic instability *in vivo* by brain microtubule associated proteins. *J. Cell Sci.* 108: 1679-1689.
8. Vandecandelaere, A., Pedrotti, B., Utton, M.A., Calvert, R.A. and Bayley, P.M. 1996. Differences in the regulation of microtubule dynamics by microtubule-associated proteins MAP-1B and MAP-2. *Cell Motil. Cytoskeleton* 35: 134-146.

CHROMOSOMAL LOCATION

Genetic locus: KRT10 (human) mapping to 17q21.2; Krt10 (mouse) mapping to 11 D.

SOURCE

MAP-1B (AA9) is a mouse monoclonal antibody raised against taxol-stabilized microtubules derived from brain of rat origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

MAP-1B (AA9) is recommended for detection of MAP-1B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for MAP-1B siRNA (h): sc-35851, MAP-1B siRNA (m): sc-35852, MAP-1B shRNA Plasmid (h): sc-35851-SH, MAP-1B shRNA Plasmid (m): sc-35852-SH, MAP-1B shRNA (h) Lentiviral Particles: sc-35851-V and MAP-1B shRNA (m) Lentiviral Particles: sc-35852-V.

Molecular Weight (predicted) of MAP-1B heavy chain: 271 kDa.

Molecular Weight (observed) of MAP-1B heavy chain: 325 kDa.


Molecular Weight of MAP-1B light chain: 34 kDa.

Positive Controls: PC-12 cell lysate: sc-2250, rat brain extract: sc-2392 or rat cerebellum extract: sc-2398.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



MAP-1B (AA9): sc-80014. Western blot analysis of MAP-1B expression in PC-12 whole cell lysate (A) and rat cerebellum tissue extract (B).



MAP-1B (AA9): sc-80014. Western blot analysis of MAP-1B expression in rat brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.