



## JAM-B (RJ-4): sc-80132

### BACKGROUND

Junctional adhesion molecule (JAM) is a member of the immunoglobulin superfamily expressed in tight junctions of epithelial cells and endothelial cells. It is implicated in transendothelial migration of leukocytes. JAM is constitutively expressed on circulating monocytes, neutrophils, lymphocytes subsets and platelets. The JAM family consists of JAM-A, JAM-B and JAM-C, alternatively designated JAM-1, JAM-2 and JAM-3, respectively. JAM-A localizes with F-Actin at the cell-cell contacts and at the membrane ruffles. It is involved in cell to cell adhesion through homophilic interactions and plays a role in the organization of tight junctions and modulation of leukocyte extravasation. JAM-B interacts with discrete subsets of PBLs, suggesting that it may play a role in lymphocyte trafficking. JAM-B and JAM-C proteins are binding partners; JAM-C may be a functional JAM-B receptor. Specifically, JAM-B adheres to T cells through heterotypic interactions with JAM-C. The JAM-B/JAM-C interaction may play a role in T, NK and dendritic cellular inflammation.

### REFERENCES

- Martin-Padura, I., et al. 1998. Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *J. Cell Biol.* 142: 117-127.
- Ozaki, H., et al. 1999. Cutting edge: combined treatment of TNF $\alpha$  and IFN- $\gamma$  causes redistribution of junctional adhesion molecule in human endothelial cells. *J. Immunol.* 163: 553-557.
- Ozaki, H., et al. 2000. Junctional adhesion molecule (JAM) is phosphorylated by protein kinase C upon platelet activation. *Biochem. Biophys. Res. Commun.* 276: 873-878.
- Ebnet, K., et al. 2000. Junctional adhesion molecule interacts with the PDZ domain-containing proteins AF-6 and ZO-1. *J. Biol. Chem.* 275: 27979-27988.
- Dejana, E., et al. 2000. The molecular organization of endothelial junctions and their functional role in vascular morphogenesis and permeability. *Int. J. Dev. Biol.* 44: 743-748.
- Bazzoni, G., et al. 2000. Homophilic interaction of junctional adhesion molecule. *J. Biol. Chem.* 275: 30970-30976.
- Arrate, M.P., et al. 2001. Cloning of human junctional adhesion molecule 3 (JAM-3) and its identification as the JAM-2 counter-receptor. *J. Biol. Chem.* 276: 45826-45832.
- Liang, T.W., et al. 2002. Vascular endothelial-junctional adhesion molecule (VE-JAM)/JAM-2 interacts with T, NK, and dendritic cells through JAM-3. *J. Immunol.* 168: 1618-1626.
- Johnson-Leger, C.A., et al. 2002. Junctional adhesion molecule-2 (JAM-2) promotes lymphocyte transendothelial migration. *Blood* 100: 2479-2486.

### CHROMOSOMAL LOCATION

Genetic locus: Jam2 (mouse) mapping to 16 C3.3.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### SOURCE

JAM-B (RJ-4) is a rat monoclonal antibody raised against full length recombinant JAM-B of mouse origin.

### PRODUCT

Each vial contains 100  $\mu$ g IgG<sub>2a</sub> in 1.0 ml PBS with < 0.1% sodium azide and protein stabilizer.

### APPLICATIONS

JAM-B (RJ-4) is recommended for detection of JAM-B of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for JAM-B siRNA (m): sc-43142, JAM-B shRNA Plasmid (m): sc-43142-SH and JAM-B shRNA (m) Lentiviral Particles: sc-43142-V.

Molecular Weight of JAM-B: 54 kDa.

Positive Controls: WRL19L cell lysate: sc-3805.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rat IgG-HRP: sc-2006 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-rat IgG-HRP: sc-2032 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.