

Cytokeratin 5/8 (C50): sc-8021

BACKGROUND

Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation which is directly applicable to the characterization of malignant tumors. Cytokeratin 5 is expressed in normal basal cells. Mutations of the Cytokeratin 5 gene (KRT5) have been shown to result in the autosomal dominant disorder epidermolysis bullosa (EB). Cytokeratin 8 expression is seen in epithelium and epithelium-derived tumors. Cytokeratins 10 and 13 are expressed highly in a subset of squamous cell carcinomas while Cytokeratin 18 is expressed in a majority of adenocarcinomas and basal cell carcinomas.

CHROMOSOMAL LOCATION

Genetic locus: KRT5/KRT8 (human) mapping to 12q13.13.

SOURCE

Cytokeratin 5/8 (C50) is a mouse monoclonal antibody raised against cytoskeletal preparation of HeLa cells.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Cytokeratin 5/8 (C50) is available conjugated to either phycoerythrin (sc-8021 PE) or Alexa Fluor® 488 (sc-8021 AF488) or Alexa Fluor® 647 (sc-8021 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Cytokeratin 5/8 (C50) is recommended for detection of Cytokeratin 5/8 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500), flow cytometry (1 µg per 1 x 10⁶ cells) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with Cytokeratin 18.

Molecular Weight of Cytokeratin 5: 58 kDa.


Molecular Weight of Cytokeratin 8: 55 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, A549 cell lysate: sc-2413 or Hep G2 cell lysate: sc-2227.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Cytokeratin 5/8 (C50): sc-8021. Western blot analysis of Cytokeratin 5/8 expression in Hep G2 (A), MCF7 (B), T24 (C), JAR (D) and A549 (E) whole cell lysates.



Cytokeratin 5/8 (C50): sc-8021. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human colon carcinoma tissue showing cytoskeletal localization (A). Cytokeratin 5/8 (C50) Alexa Fluor® 488: sc-8021 AF488 Direct immunofluorescence staining of formalin-fixed SW480 cells showing cytoskeletal localization. Blocked with UltraCruz® Blocking Reagent: sc-516214 (B).

SELECT PRODUCT CITATIONS

- Kim, H.L., et al. 2001. Mitogen-activated protein kinase kinase 4 metastasis suppressor gene expression is inversely related to histological pattern in advancing human prostatic cancers. *Cancer Res.* 61: 2833-2837.
- Pearce, V.P., et al. 2008. Immortalization of epithelial progenitor cells mediated by resveratrol. *Oncogene* 27: 2365-2374.
- Carmona, F.D., et al. 2010. Development of the cornea of true moles (*Talpidae*): morphogenesis and expression of PAX6 and cytokeratins. *J. Anat.* 217: 488-500.
- Wei, Q., et al. 2013. Keratinocyte cytoskeletal roles in cell sheet engineering. *BMC Biotechnol.* 13: 17.
- Iannolo, G., et al. 2016. Numb expression contributes to the maintenance of an undifferentiated state in human epidermis. *Cell Transplant.* 25: 353-364.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA