

Ox40L (RR19): sc-80271

BACKGROUND

The Ox40 ligand, Ox40L (also designated gp34), is a type II membrane protein, due to the absence of a signal peptide. Ox40L, a member of the tumor necrosis factor (TNF) superfamily, is a costimulatory molecule involved in dendritic cell-T cell interactions, T cell homing and B cell activation. Engagement of Ox40L with its receptor, Ox40, delivers a strong costimulatory signal to effector T cells. Ox40L is found preferentially on activated B cells and its receptor, Ox40, is a member of the tumor necrosis factor receptor (TNFR) family that is expressed on activated T cells. Ox40L plays a critical role in antigen-specific T cell responses *in vivo* and in both the priming and effector phases of T cell activation when expressed on antigen-presenting cells (APCs).

REFERENCES

- Hahne, M., Kataoka, T., Schroter, M., Hofman, K., Irmler, M., Bodmer, J.L., Schneider, P., Bornand, T., Holler, N., French, L.E., Sordat, B., Rimoldi, D. and Tschoopp, J. 1998. APRIL, a new ligand of the tumor necrosis factor family, stimulates tumor cell growth. *J. Exp. Med.* 188: 1185-1190.
- Chen, A.I., McAdam, A.J., Buhlmann, J.E., Scott, S., Luper, M.L., Jr., Greenfield, E.A., Baum, P.R., Fanslow, W.C., Calderhead, D.M., Freeman, G.J. and Sharpe, A.H. 1999. Ox40-ligand has a critical costimulatory role in dendritic cell-T cell interactions. *Immunity* 11: 689-698.
- Matsumura, Y., Hori, T., Kawamata, S., Imura, A. and Uchiyama, T. 1999. Intracellular signaling of gp34, the Ox40 ligand: induction of c-Jun and c-Fos mRNA expression through gp34 upon binding of its receptor, Ox40. *J. Immunol.* 163: 3007-3011.
- Weinberg, A.D., Rivera, M.M., Prell, R., Morris, A., Ramstad, T., Vetto, J.T., Urba, W.J., Alvord, G., Bunce, C. and Shields, J. 2000. Engagement of the Ox40 receptor *in vivo* enhances antitumor immunity. *J. Immunol.* 164: 2160-2169.
- Morimoto, S., Kanno, Y., Tanaka, Y., Tokano, Y., Hashimoto, H., Jacquot, S., Morimoto, C., Schlossman, S.F., Yagita, H., Okumura, K. and Kobata, T. 2000. CD134L engagement enhances human B cell Ig production: CD154/CD40, CD70/CD27 and CD134/CD134L interactions coordinately regulate T cell-dependent B cell responses. *J. Immunol.* 164: 4097-4104.
- Murata, K., Ishii, N., Takano, H., Miura, S., Ndhlovu, L.C., Nose, M., Noda, T. and Sugamura, K. 2000. Impairment of antigen-presenting cell function in mice lacking expression of Ox40 ligand. *J. Exp. Med.* 191: 365-374.
- Malmstrom, V., Shipton, D., Singh, B., Al-Shamkhani, A., Puklavec, M.J., Barclay, A.N. and Powrie, F. 2001. CD134L expression on dendritic cells in the mesenteric lymph nodes drives colitis in T cell-restored SCID mice. *J. Immunol.* 166: 6972-6981.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: TNFSF4 (human) mapping to 1q25.1.

SOURCE

Ox40L (RR19) is a mouse monoclonal antibody raised against the extracellular domain of Ox40L of human origin.

PRODUCT

Each vial contains 100 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and protein stabilizer. Also available azide-free for neutralization, sc-80271 L, 100 µg/0.1 ml.

APPLICATIONS

Ox40L (RR19) is recommended for detection of Ox40L of human origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 µg per 1 × 10⁶ cells).

Suitable for use as control antibody for Ox40L siRNA (h): sc-42824, Ox40L shRNA Plasmid (h): sc-42824-SH and Ox40L shRNA (h) Lentiviral Particles: sc-42824-V.

Molecular Weight of Ox40L: 22 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Immunofluorescence: use m-IgG_κ BP-FITC: sc-516140 or m-IgG_κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

SELECT PRODUCT CITATIONS

- Ilves, T. and Harvima, I.T. 2013. Ox40 ligand and Ox40 are increased in atopic dermatitis lesions but do not correlate with clinical severity. *J. Eur. Acad. Dermatol. Venereol.* 27: e197-e205.
- Zhang, J.Y., Liu, B., Wang, Y.N., Zhang, W.N. and Wang, F.J. 2014. Effect of rosuvastatin on Ox40L and PPAR-γ expression in human umbilical vein endothelial cells and atherosclerotic cerebral infarction patients. *J. Mol. Neurosci.* 52: 261-268.

RESEARCH USE

For research use only, not for use in diagnostic procedures.