



Thioguanine (5B2): sc-80593

BACKGROUND

Thioguanine is an antineoplastic compound that is clinically used in the treatment of cancer, mainly acute leukemias and chronic myeloid leukemias. Thioguanine is a guanine analog with a molecular weight of 167.19 g/mol and a half life ranging from 25 to 240 minutes with an average of about 80 minutes. Thioguanine has antimetabolite action. Once inside the cell, Thioguanine is transformed into 6-thioguanosine 5'-monophosphate (TGMP) which functions by interfering with the synthesis of guanine nucleotides by pseudo-feedback interference with purine biosynthesis. This results in the induction of cell cycle arrest and apoptosis. Some of the activity of Thioguanine may also be due to the incorporation of Thioguanine nucleotides into both RNA and DNA.

REFERENCES

1. Chojnacki, H. and Sokalski, W.A. 1975. Interactions of 6-Thioguanine in B-DNA: possible mechanism of its mutagenic action. *J. Theor. Biol.* 54: 167-174.
2. The Finnish Leukaemia Group. 1979. The effect of Thioguanine on a combination of daunorubicine, cytarabine and prednisone in the treatment of acute leukaemia in adults. *Scand. J. Haematol.* 23: 124-128.
3. Dooley, T. and Maddocks, J.L. 1982. Assay of an active metabolite of 6-Thioguanine, 6-thioguanosine 5'-monophosphate, in human red blood cells. *J. Chromatogr.* 229: 121-127.
4. Cleaver, J.E. 1984. Differential toxicity of 3-Aminobenzamide to wildtype and 6-Thioguanine-resistant Chinese hamster cells by interference with pathways of purine biosynthesis. *Mutat. Res.* 131: 123-127.
5. Sriram, S. 1994. Longitudinal study of frequency of HPRT mutant T cells in patients with multiple sclerosis. *Neurology* 44: 311-315.
6. Krynetski, E.Y., Krynetskaia, N.F., Yanishevski, Y. and Evans, W.E.. 1995. Methylation of mercaptopurine, Thioguanine, and their nucleotide metabolites by heterologously expressed human thiopurine S-methyltransferase. *Mol. Pharmacol.* 47: 1141-1147.
7. Grant, D.D., Goldstein, R., Karsh, J. and Birnboim, H.C. 1999. Elimination of non-viable 6-Thioguanine-sensitive T cells from viable T cells prior to PCR analysis. *J. Immunol. Methods* 225: 61-66.
8. Bohon, J. and de los Santos, C.R. 2005. Effect of 6-Thioguanine on the stability of duplex DNA. *Nucleic Acids Res.* 33: 2880-2886.
9. Parrish, J.A. 2005. Immunosuppression, skin cancer, and ultraviolet A radiation. *N. Engl. J. Med.* 353: 2712-2713.

SOURCE

Thioguanine (5B2) is a mouse monoclonal antibody raised against 9-substituted Thioguanine linked to PPD.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 100 µg IgG_{2a} in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Thioguanine (5B2) is recommended for detection of Thioguanine by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with a variety of Thioguanine analogues; non cross-reactive with 6-(methylthio)purine, 6-thioxanthine, 2-amino-8-phenyl-6-purinethione and 6-(methylthio)purine riboside.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.