

Pepsin A (8003): sc-80914

BACKGROUND

Pepsin is one of the main proteolytic enzymes secreted by the gastric mucosa. Pepsin consists of a single polypeptide chain and arises from its precursor, pepsinogen, by removal of a 41 amino acid segment from the N-terminus. Pepsinogen is synthesized in the stomach lining, and hydrochloric acid, also produced by the gastric mucosa, is necessary to convert the inactive enzyme and to maintain the optimum acidity (pH 1-3) for Pepsin function. Pepsin is particularly effective in cleaving peptide bonds involving aromatic amino acids. Pepsin shows extremely broad specificity; although bonds involving phenylalanine and leucine are preferred, many others are also cleaved to some extent. Pepsin A is a member of the subfamily A1 within the Pepsin family and is the predominant endopeptidase in the gastric juice of vertebrates. Pepsin A is inhibited by ovUS-1, a uterine Serpin.

REFERENCES

- Hampson, I.N., Hampson, L., Pinkoski, M., Cross, M., Heyworth, C.M., Bleackley, R.C., Atkinson, E. and Dexter, T.M. 1997. Identification of a serpin specifically expressed in multipotent and bipotent hematopoietic progenitor cells and in activated T cells. *Blood* 89: 108-118.
- Hamerman, J.A., Hayashi, F., Schroeder, L.A., Gygi, S.P., Haas, A.L., Hampson, L., Coughlin, P., Aebbersold, R. and Aderem, A. 2002. Serpin 2A is induced in activated macrophages and conjugates to a ubiquitin homolog. *J. Immunol.* 168: 2415-2423.
- Hampson, L., Hampson, I.N., Babichuk, C.K., Cotter, L., Bleackley, R.C., Dexter, T.M. and Cross, M.A. 2002. A minimal serpin promoter with high activity in haematopoietic progenitors and activated T cells. *Hematol. J.* 2: 150-160.
- Gettins, P.G. 2002. Serpin structure, mechanism, and function. *Chem. Rev.* 102: 4751-4804.
- Carrell, R.W. and Huntington, J.A. 2003. How serpins change their fold for better and for worse. *Biochem. Soc. Symp.* 70: 163-178.
- Morris, E.C., Dafforn, T.R., Forsyth, S.L., Missen, M.A., Horvath, A.J., Hampson, L., Hampson, I.N., Currie, G., Carrell, R.W. and Coughlin, P.B. 2003. Murine Serpin 2A is a redox-sensitive intracellular protein. *Biochem. J.* 371: 165-173.
- van Gent, D., Sharp, P., Morgan, K. and Kalsheker, N. 2003. Serpins: structure, function and molecular evolution. *Int. J. Biochem. Cell Biol.* 35: 1536-1547.
- Stanley, P. and Stein, P.E. 2003. BmSPN2, a serpin secreted by the filarial nematode *Brugia malayi*, does not inhibit human neutrophil proteinases but plays a noninhibitory role. *Biochemistry* 42: 6241-6248.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: PGA3/PGA4/PGA5 (human) mapping to 11q12.2.

SOURCE

Pepsin A (8003) is a mouse monoclonal antibody raised against purified Pepsinogen A of human origin.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Pepsin A (8003) is recommended for detection of Pepsinogen A and Pepsin A of human origin by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Pepsin A siRNA (h): sc-61317, Pepsin A shRNA Plasmid (h): sc-61317-SH and Pepsin A shRNA (h) Lentiviral Particles: sc-61317-V.

Molecular Weight of Pepsin A: 42 kDa.

RESEARCH USE

For research use only, not for use in diagnostic procedures.