

SERCA2 (C-20): sc-8094

BACKGROUND

ATP dependent calcium pumps are responsible, in part, for the maintenance of low cytoplasmic free calcium concentrations. The ATP pumps that reside in intracellular organelles are encoded by a family of structurally related enzymes, termed the sarcoplasmic or endoplasmic reticulum calcium (SERCA) ATPases. The sarcoplasmic reticulum of striated muscle is a specialized intracellular membrane system that plays a critical role in the contraction and relaxation of muscle. The SERCAs mediate Ca^{2+} uptake into intracellular stores. SERCA-mediated Ca^{2+} uptake induces and maintains muscular relaxation. The SERCA1 gene is exclusively expressed in type II (fast) skeletal muscle. The SERCA2 gene is subject to tissue-dependent processing which is responsible for the generation of the SERCA2a muscle-specific form expressed in type I (slow) skeletal, cardiac and smooth muscle, and the SERCA2b isoform expressed in all cell types. The SERCA3 gene is not as well characterized and is found in non-muscle cells. SERCA2 plays an important part in regulating cardiac contractile function. SERCA3 is an isoform expressed in several cell types including platelets, lymphoid cells and mast cells. SERCA1, SERCA2 and SERCA3 all undergo alternative splicing.

CHROMOSOMAL LOCATION

Genetic locus: ATP2A2 (human) mapping to 12q24.11; Atp2a2 (mouse) mapping to 5 F.

SOURCE

SERCA2 (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of SERCA2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-8094 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SERCA2 (C-20) is recommended for detection of SERCA2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SERCA2 (C-20) is also recommended for detection of SERCA2 in additional species, including canine, porcine, avian and feline.

Suitable for use as control antibody for SERCA2 siRNA (h): sc-36484, SERCA2 siRNA (m): sc-36485, SERCA2 shRNA Plasmid (h): sc-36484-SH, SERCA2 shRNA Plasmid (m): sc-36485-SH, SERCA2 shRNA (h) Lentiviral Particles: sc-36484-V and SERCA2 shRNA (m) Lentiviral Particles: sc-36485-V.


Molecular Weight of SERCA2: 100 kDa.

Positive Controls: rat heart extract: sc-2393, T98G cell lysate: sc-2294 or rat cardiac tissue extract.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



SERCA2 (C-20): sc-8094. Western blot analysis of SERCA2 expression in rat heart extract.

SELECT PRODUCT CITATIONS

1. Beutner, G., et al. 2001. Identification of a ryanodine receptor in rat heart mitochondria. *J. Biol. Chem.* 276: 21482-21488.
2. Tishkoff, D.X., et al. 2008. Functional vitamin D receptor (VDR) in the t-tubules of cardiac myocytes: VDR knockout cardiomyocyte contractility. *Endocrinology* 149: 558-564.
3. Zhao, G., et al. 2010. Interaction between vitamin D receptor with caveolin-3 and regulation by 1,25-dihydroxyvitamin D₃ in adult rat cardiomyocytes. *J. Steroid Biochem. Mol. Biol.* 121: 159-163.
4. Lin, H.H., et al. 2010. Neuronatin promotes neural lineage in ESCs via Ca^{2+} signaling. *Stem Cells* 28: 1950-1960.
5. Rottlaender, D., et al. 2010. Connexin 43 acts as a cytoprotective mediator of signal transduction by stimulating mitochondrial KATP channels in mouse cardiomyocytes. *J. Clin. Invest.* 120: 1441-1453.
6. McMullen, D.C., et al. 2010. In cold-hardy insects, seasonal, temperature, and reversible phosphorylation controls regulate sarco/endoplasmic reticulum Ca^{2+} -ATPase (SERCA). *Physiol. Biochem. Zool.* 83: 677-686.
7. Burmeister, D., et al. 2012. Impact of partial urethral obstruction on bladder function: time-dependent changes and functional correlates of altered expression of Ca^{2+} signaling regulators. *Am. J. Physiol. Renal Physiol.* 302: F1517-F1528.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
Satisfation
Guaranteed

Try **SERCA2 (F-1): sc-376235** or **SERCA2 (IID8): sc-53010**, our highly recommended monoclonal alternatives to SERCA2 (C-20). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **SERCA2 (F-1): sc-376235**.