

PECAM-1 (2F7B2): sc-81158

BACKGROUND

Cell adhesion molecules are a family of closely related cell surface glycoproteins involved in cell-cell interactions during growth and are thought to play an important role in embryogenesis and development. Neuronal cell adhesion molecule (NCAM) expression is observed in a variety of human tumors including neuroblastomas, rhabdomyosarcomas, Wilms' tumors, Ewing's sarcomas and some primitive myeloid malignancies. The intracellular adhesion molecule-1 (ICAM-1), also referred to as CD54, is an integral membrane protein of the immunoglobulin superfamily and recognizes the $\beta 2/\alpha 1$ and $\beta 2/\alpha M$ Integrins. PECAM-1 (platelet/endothelial cell adhesion molecule-1), also referred to as CD31, is a glycoprotein expressed on the cell surfaces of monocytes, neutrophils, platelets and a subpopulation of T cells. VCAM-1 (vascular cell adhesion molecule-1) was first identified as an adhesion molecule induced on human endothelial cells by inflammatory cytokines such as IL-1, tumor necrosis factor (TNF) and lipopolysaccharide (LPS). The KALIG gene encodes a nerve cell adhesion molecule (NCAM)-like protein and is deleted in 66% of patients with Kallmann's syndrome, anosmia with secondary hypogonadism.

REFERENCES

1. Patel, K., et al. 1993. Vase mini-exon usage by NCAM is not restricted to tumours of neuroectodermal origin. *Int. J. Cancer* 54: 772-777.
2. Cowen, M.A. and Green, M. 1993. The Kallmann's syndrome variant (KSV) model of the schizophrenias. *Schizophr. Res.* 9: 1-10.

CHROMOSOMAL LOCATION

Genetic locus: PECAM1 (human) mapping to 17q23.3.

SOURCE

PECAM-1 (2F7B2) is a mouse monoclonal antibody raised against a recombinant protein corresponding to amino acids 621-738 of PECAM-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PECAM-1 (2F7B2) is recommended for detection of PECAM-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for PECAM-1 siRNA (h): sc-29445, PECAM-1 shRNA Plasmid (h): sc-29445-SH and PECAM-1 shRNA (h) Lentiviral Particles: sc-29445-V.


Molecular Weight of PECAM-1: 130 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, MOLT-4 cell lysate: sc-2233 or THP-1 cell lysate: sc-2238.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



PECAM-1 (2F7B2): sc-81158. Western blot analysis of PECAM-1 expression in THP-1 (A), CCRF-CEM (B), Jurkat (C), MOLT-4 (D) and IMR-32 (E) whole cell lysates and human lung tissue extract (F).



PECAM-1 (2F7B2): sc-81158. Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing membrane staining of endothelial cells.

SELECT PRODUCT CITATIONS

1. Zhang, J., et al. 2017. Palmitate impairs angiogenesis via suppression of cathepsin activity. *Mol. Med. Rep.* 15: 3644-3650.
2. Duan, F., et al. 2018. Biphasic modulation of Insulin signaling enables highly efficient hematopoietic differentiation from human pluripotent stem cells. *Stem Cell Res. Ther.* 9: 205.
3. Saadane, A., et al. 2019. Retinal vascular abnormalities and microglia activation in mice with deficiency in cytochrome P450 46A1-mediated cholesterol removal. *Am. J. Pathol.* 189: 405-425.
4. Li, X., et al. 2022. Type II collagen-positive progenitors are important stem cells in controlling skeletal development and vascular formation. *Bone Res.* 10: 46.
5. Dong, R., et al. 2023. Rutin alleviates EndMT by restoring autophagy through inhibiting HDAC1 via PI3K/AKT/mTOR pathway in diabetic kidney disease. *Phytomedicine* 112: 154700.
6. Li, M., et al. 2024. Human pluripotent stem cells derived endothelial cells repair choroidal ischemia. *Adv. Sci.* 11: e2302940.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **PECAM-1 (H-3): sc-376764** for PECAM-1 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.