

BAT1/DDX39 (2060C10a): sc-81196

BACKGROUND

BAT1 (spliceosome RNA helicase BAT1), also known as DEAD-box protein UAP56 (56 kDa U2AF65-associated protein), HLA-B associated transcript-1 or ATP-dependent RNA helicase p47, is a member of the DECD subfamily of DEAD-box helicases. Important for mRNA splicing and nuclear export, BAT1 interacts with the mRNA export factor ALY and also functions as a splicing factor, mediating the first ATP-dependent step of spliceosome assembly. BAT1 associates with transcription elongation factor THO proteins, forming the TREX complex, and also interacts with splicing machinery to form the exon junction complex. Due to alternative splicing events, BAT1 exists in two isoforms. DDX39 (DEAD box protein 39), like BAT1, is a member of the DEAD-box family of helicases. Localized to the nucleus and expressed in lung, brain, kidney, spleen, thymus and salivary gland, DDX39 functions in a similar manner to BAT1 and is involved in pre-mRNA splicing and mRNA export out of the nucleus. DDX39 expression is upregulated in lung squamous cell carcinoma, suggesting a role for DDX39 in tumorigenesis.

REFERENCES

- Fleckner, J., Zhang, M., Valcárcel, J. and Green, M.R. 1997. U2AF65 recruits a novel human DEAD box protein required for the U2 snRNP-branchpoint interaction. *Genes Dev.* 11: 1864-1872.
- Momose, F., Basler, C.F., O'Neill, R.E., Iwamatsu, A., Palese, P. and Nagata, K. 2001. Cellular splicing factor RAF-2p48/NPI-5/BAT1/UAP56 interacts with the influenza virus nucleoprotein and enhances viral RNA synthesis. *J. Virol.* 75: 1899-1908.
- Luo, M.L., Zhou, Z., Magni, K., Christoforides, C., Rappsilber, J., Mann, M. and Reed, R. 2001. Pre-mRNA splicing and mRNA export linked by direct interactions between UAP56 and Aly. *Nature* 413: 644-647.
- Price, P., Wong, A.M., Williamson, D., Voon, D., Baltic, S., Allcock, R.J., Boodhoo, A. and Christiansen, F.T. 2004. Polymorphisms at positions -22 and -348 in the promoter of the BAT1 gene affect transcription and the binding of nuclear factors. *Hum. Mol. Genet.* 13: 967-974.
- Shi, H., Cordin, O., Minder, C.M., Linder, P. and Xu, R.M. 2004. Crystal structure of the human ATP-dependent splicing and export factor UAP56. *Proc. Natl. Acad. Sci. USA* 101: 17628-17633.
- Thakurta, A.G., Gopal, G., Yoon, J.H., Kozak, L. and Dhar, R. 2005. Homolog of BRCA2-interacting Dss1p and UA56p link Mlo3p and Rae1p for mRNA export in fission yeast. *EMBO J.* 24: 2512-2523.
- Kapadia, F., Pryor, A., Chang, T.H. and Johnson, L.F. 2006. Nuclear localization of polyA⁺ mRNA following siRNA reduction of expression of the mammalian RNA helicases UAP56 and URH49. *Gene* 384: 37-44.
- Ramasawmy, R., Cunha-Neto, E., Faé, K.C., Müller, N.G., Cavalcanti, V.L., Drigo, S.A., Ianni, B., Mady, C., Kalil, J. and Goldberg, A.C. 2006. BAT1, a putative anti-inflammatory gene, is associated with chronic Chagas cardiomyopathy. *J. Infect. Dis.* 193: 1394-1399.

CHROMOSOMAL LOCATION

Genetic locus: BAT1 (human) mapping to 6p21.33, DDX39 (human) mapping to 19p13.12.

SOURCE

BAT1/DDX39 (2060C10a) is a mouse monoclonal antibody raised against a recombinant protein corresponding to the C-terminal region of BAT1 of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% BSA.

APPLICATIONS

BAT1/DDX39 (2060C10a) is recommended for detection of BAT1 and DDX39 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].


Molecular Weight of BAT1: 48 kDa.

Molecular Weight of DDX39: 49 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



BAT1/DDX39 (2060C10a): sc-81196. Western Blot analysis of human recombinant BAT1 fusion protein.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.