SANTA CRUZ BIOTECHNOLOGY, INC.

Dmp1 (DMTF5I250): sc-81249



BACKGROUND

The highly leukemogenic avian retrovirus E26 contains two oncogenes, v-Myb and v-Ets, which are expressed together as a fusion protein. The cellular homolog of v-Myb, designated c-Myb, encodes a transcription factor. Deletion or disruption of a negative regulatory domain mapping within the carboxy terminus of c-Myb results in enhanced transactivating capacity and in parallel, leads to activation of its ability to transform hemopoietic cells. c-Myb is expressed preferentially, but not exclusively, in immature hemopoietic cells and its expression decreases as cells differentiate. A second member of the Myb proto-oncogene family, B-Myb, encodes another sequence-specific DNA binding protein. Studies suggest that B-Myb expression rescues cells from p53-induced G_1 arrest mediated by p21. Dmp1 (also designated cyclin D binding Myb-like transcription factor 1) has also been identified as a Myb-like transcription factor. It contains three tandem Myb repeats and has been shown to be a substrate for cyclin D-dependent kinases.

REFERENCES

- Gonda, T.J., et al. 1984. Expression of Myb, Myc and Fos proto-oncogenes during the differentiation of a murine myeloid leukaemia. Nature 310: 249-251.
- Gonda, T.J., et al. 1985. Nucleotide sequence of cDNA clones of the murine Myb proto-oncogene. EMBO J. 4: 2004-2008.
- Sakura, H., et al. 1989. Delineation of three functional domains of the transcriptional activator encoded by the c-Myb protooncogene. Proc. Natl. Acad. Sci. USA 86: 5758-5762.
- Mizuguchi, G., et al. 1990. DNA binding activity and transcriptional activator function of the human B-Myb protein compared with c-Myb. J. Biol. Chem. 265: 9280-9284.
- Ramsay, R.G., et al. 1991. Increase in specific DNA binding by carboxyl truncation suggests a mechanism for activation of Myb. Oncogene 6: 1875-1879.
- Favier, D., et al. 1994. Detection of proteins that bind to the leucine zipper motif of c-Myb. Oncogene 9: 305-311.
- Lin, D., et al. 1994. Constitutive expression of B-Myb can bypass p53induced Waf1/Cip1-mediated G₁ arrest. Proc. Natl. Acad. Sci. USA 91: 10079-10083.
- Mallakin, A., et al. 2006. Expression of Dmp1 in specific differentiated, nonproliferating cells and its regulation by E2Fs. Oncogene 25: 7703-7713.
- 9. Inoue, K., et al. 2007. Dmp1 and tumor suppression. Oncogene 26: 4329-4335.

CHROMOSOMAL LOCATION

Genetic locus: DMTF1 (human) mapping to 7q21.12.

STORAGE

For immediate and continuous use, store at 4° C for up to one month. For sporadic use, freeze in working aliquots in order to avoid repeated freeze/ thaw cycles. If turbidity is evident upon prolonged storage, clarify solution by centrifugation.

SOURCE

Dmp1 (DMTF51250) is a mouse monoclonal antibody raised against a recombinant protein fragment corresponding to an internal region of Dmp1 of human origin.

PRODUCT

Each vial contains 100 $\mu g~lg G_1$ kappa light chain in 1.0 ml of PBS with <0.1% sodium azide and 1.0% stabilizer protein.

APPLICATIONS

Dmp1 (DMTF5I250) is recommended for detection of Dmp1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for Dmp1 siRNA (h): sc-38068, Dmp1 shRNA Plasmid (h): sc-38068-SH and Dmp1 shRNA (h) Lentiviral Particles: sc-38068-V.

Molecular Weight of Dmp1: 84 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).





Dmp1 (DMTF5I250): sc-81249. Western Blot analysis of human recombinant Dmp1 fusion protein.

SELECT PRODUCT CITATIONS

- Wang, Y., et al. 2022. MicroRNA-196a-5p overexpression in Wharton's jelly umbilical cord stem cells promotes their osteogenic differentiation and new bone formation in bone defects in the rat calvarium. Cell Tissue Res. 390: 245-260.
- 2. Deng, P., et al. 2023. The role of EMILIN-1 in the osteo/odontogenic differentiation of dental pulp stem cells. BMC Oral Health 23: 203.

RESEARCH USE

For research use only, not for use in diagnostic procedures.