

E2F-1 (658C2a): sc-81257

BACKGROUND

The human retinoblastoma gene product appears to play an important role in the negative regulation of cell proliferation. Functional inactivation of Rb can be mediated either through mutation or as a consequence of interaction with DNA tumor virus encoded proteins. Of all the Rb associations described to date, the identification of a complex between Rb and the transcription factor E2F most directly implicates Rb in regulation of cell proliferation. E2F was originally identified through its role in transcriptional activation of the adenovirus E2 promoter. Sequences homologous to the E2F binding site have been found upstream of a number of genes that encode proteins with putative functions in the G₁ and S phases of the cell cycle. E2F-1 is a member of a broader family of transcriptional regulators including E2F-2, E2F-3, E2F-4, E2F-5 and E2F-6, each of which forms heterodimers with a second protein, DP-1, forming an "active" E2F transcriptional regulatory complex.

REFERENCES

- Chellappan, S., et al. 1991. The E2F transcription factor is a cellular target for the Rb protein. *Cell* 65: 1053-1061.
- Chittenden, T., et al. 1991. The T/E1A-binding domain of the retinoblastoma product can interact selectively with a sequence-specific DNA-binding protein. *Cell* 65: 1073-1082.
- Helin, K., et al. 1992. A cDNA encoding a pRb-binding protein with properties of the transcription factor E2F. *Cell* 70: 337-350.
- Helin, K., et al. 1993. Heterodimerization of the transcription factors E2F-1 and DP-1 leads to cooperative transactivation. *Genes Dev.* 7: 1850-1861.
- Krek, W., et al. 1993. Binding to DNA and the retinoblastoma gene product promoted by complex formation of different E2F family members. *Science* 262: 1557-1560.
- Ginsberg, D., et al. 1994. E2F-4, a new member of the E2F transcription factor family, interacts with p107. *Genes Dev.* 8: 2665-2679.
- Beijersbergen, R.L., et al. 1994. E2F-4, a new member of the E2F gene family, has oncogenic activity and associates with p107 *in vivo*. *Genes Dev.* 8: 2680-2690.
- Trimarchi, J.M., et al. 1998. E2F-6, a member of the E2F family that can behave as a transcriptional repressor. *Proc. Natl. Acad. Sci. USA* 95: 2850-2855.

CHROMOSOMAL LOCATION

Genetic locus: E2F1 (human) mapping to 20q11.22.

SOURCE

E2F-1 (658C2a) is a mouse monoclonal antibody raised against a recombinant protein corresponding to an internal region of E2F-1 of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% BSA.

APPLICATIONS

E2F-1 (658C2a) is recommended for detection of E2F-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for E2F-1 siRNA (h): sc-29297, E2F-1 shRNA Plasmid (h): sc-29297-SH and E2F-1 shRNA (h) Lentiviral Particles: sc-29297-V.


Molecular Weight of E2F-1: 60 kDa.

Positive Controls: E2F-1 (h): 293T lysate: sc-116137, K-562 nuclear extract: sc-2130 or MOLT-4 cell lysate: sc-2233.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



E2F-1 (658C2a): sc-81257. Western blot analysis of E2F-1 expression in non-transfected: sc-117752 (A) and human E2F-1 transfected: sc-116137 (B) 293T whole cell lysates.



E2F-1 (658C2a): sc-81257. Western blot analysis of E2F-1 expression in non-transfected: sc-117752 (A) and human E2F-1 transfected: sc-116659 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Wei, Y., et al. 2010. Regulation of the β1,4-Galactosyltransferase I promoter by E2F1. *J. Biochem.* 148: 263-271.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.