SANTA CRUZ BIOTECHNOLOGY, INC.

NAB1 (2607C2a): sc-81565



BACKGROUND

Transcriptional control is in part regulated by interactions between DNA-bound transcription factors, such as Egr-1/NGFI-A, and coregulatory proteins such as NAB (for NGFI-A-binding proteins). The evolutionarily conserved NAB proteins, NAB1 and NAB2, are corepressors of Egr-1/NGFI-A. Both NAB1 and NAB2 contain an amino-terminal NAB-conserved domain 1 (NCB1), which is required for binding NGFI-A, and a carboxy-terminal NCD2, which is responsible for the repressor function of NAB proteins. NAB1 requires NGFI-A to gain access to DNA, indicating that NAB1 is an active repressor that works by a direct mechanism. NAB1, which is constitutively expressed, is localized exclusively in the nucleus and may play a role in controlling processes such as cell division, differentiation and apoptosis.

REFERENCES

- 1. Russo, M.W., et al. 1993. Transcriptioanl activity of the zinc finger protein NGFI-A is influenced by its interaction with a cellular factor. Mol. Cell. Biol. 13: 6858-6865.
- 2. Russo, M.W., et al. 1995. Identification of NAB1, a repressor of NGFI-A- and Krox-20-mediated transcription. Proc. Natl. Acad. Sci. USA 92: 6873-6877.
- 3. Svaren, J., et al. 1996. NAB2, a corepressor of NGFI-A (Egr-1) and Krox-20, is induced by proliferative and differentiative stimuli. Mol. Cell. Biol. 16: 3545-3553.
- 4. Swirnoff, A.H., et al. 1998. NAB1, a corepressor of NGFI-A (Egr-1), contains an active transcriptional repression domain. Mol. Cell. Biol. 18: 512-524.
- 5. Sevetson, B.R., et al. 2000. A novel activation function for NAB proteins in EGR-dependent transcription of the luteinizing hormone β gene. J. Biol. Chem. 275: 9749-9757.
- 6. Braddock, M., et al. 2000. Therapeutic applications of the transcriptional corepressor proteins NAB1 and NAB2 in regenerative medicine. Drugs 3: 783-787.
- 7. Mussgnug, J.H., et al. 2005. NAB1 is an RNA binding protein involved in the light-regulated differential expression of the light-harvesting antenna of Chlamvdomonas reinhardtii. Plant Cell 17: 3409-3421.
- 8. Buitrago, M., et al. 2005. The transcriptional repressor NAB1 is a specific regulator of pathological cardiac hypertrophy. Nat. Med. 11: 837-844.
- 9. Forster, B., et al. 2006. Comparative proteomics of high light stress in the model alga Chlamydomonas reinhardtii. Proteomics 6:4309-4320.

CHROMOSOMAL LOCATION

Genetic locus: NAB1 (human) mapping to 2q32.2; Nab1 (mouse) mapping to 1 C1.1.

SOURCE

NAB1 (2607C2a) is a mouse monoclonal antibody raised against a recombinant protein corresponding to a region near the C-terminus of NAB1 of human origin.

PRODUCT

Each vial contains 100 μ g lgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 1.0% stabilizer protein.

APPLICATIONS

NAB1 (2607C2a) is recommended for detection of NAB1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for NAB1 siRNA (h): sc-38089, NAB1 siRNA (m): sc-38090, NAB1 shRNA Plasmid (h): sc-38089-SH, NAB1 shRNA Plasmid (m): sc-38090-SH, NAB1 shRNA (h) Lentiviral Particles: sc-38089-V and NAB1 shRNA (m) Lentiviral Particles: sc-38090-V.

Molecular Weight of NAB1: 54 kDa.

Positive Controls: NAB1 (m): 293T Lysate: sc-121928, NAB1 (h): 293T Lysate: sc-115360 or SK-N-MC cell lysate: sc-2237.

DATA





NAB1 (2607C2a): sc-81565. Western blot analysis of NAB1 expression in non-transfected 293T: sc-117752 (A), human NAB1 transfected 293T: sc-115360 (B) and SK-N-MC (C) whole cell lysates

NAB1 (2607C2a): sc-81565. Western blot analysis of NAB1 expression in non-transfected: sc-117752 (A) and mouse NAB1 transfected: sc-121928 (B) 293T whole cell lysates

SELECT PRODUCT CITATIONS

- 1. Ren, Y.R., et al. 2012. Unbiased discovery of interactions at a control locus driving expression of the cancer-specific therapeutic and diagnostic target, mesothelin. J. Proteome Res. 11: 5301-5310.
- 2. Hancock, M.H., et al. 2020. Human cytomegalovirus miRNAs regulate TGF- β to mediate myelosuppression while maintaining viral latency in CD34+ hematopoietic progenitor cells. Cell Host Microbe 27: 104-114.e4.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.