

# Hep B Pol (9F9): sc-81592

## BACKGROUND

The Hep B Pol (Hepatitis B polymerase) protein is an enzyme that facilitates the transcription of the Hep B RNA genome into the dsDNA found within the forming viral capsids. Hep B Pol performs multiple functions, as it possesses the ability to copy both DNA and RNA templates as well as to excise the RNA strand from a DNA-RNA heteroduplex. Hep B Pol is encapsidated by a continuous core shell together with pregenomic (pg) RNA. It is here that reverse transcription is initiated by the Hep B Pol protein binding to the epsilon loop of the pgRNA. The reaction is primed by the Hep B Pol protein covalently binding to a negative (-) sense DNA strand which is extended into a complete strand followed by synthesis of the positive + strand. The Hep B Pol protein is activated by HSP 70 and HSP 40 garnered from the host cell and can be inactivated by a variety of reverse transcriptase inhibitors. Hep B Pol contains a terminal protein domain (TP) which is specific to Hepadnaviruses.

## REFERENCES

1. Valenzuela, P., Gray, P., Quiroga, M., Zaldivar, J., Goodman, H.M. and Rutter, W.J. 1979. Nucleotide sequence of the gene coding for the major protein of hepatitis B virus surface antigen. *Nature* 280: 815-819.
2. Lai, C.L. and Yuen, M.F. 2000. Profound suppression of hepatitis B virus replication with lamivudine. *J. Med. Virol.* 61: 367-373.
3. Hadziyannis, S.J. and Papatheodoridis, G.V. 2004. Adefovir dipivoxil in the treatment of chronic hepatitis B virus infection. *Expert Rev. Anti Infect. Ther.* 2: 475-483.
4. Langley, D.R., Walsh, A.W., Baldick, C.J., Eggers, B.J., Rose, R.E., Levine, S.M., Kapur, A.J., Colonno, R.J. and Tenney, D.J. 2007. Inhibition of hepatitis B virus polymerase by entecavir. *J. Virol.* 81: 3992-4001.
5. Beck, J. and Nassal, M. 2007. Hepatitis B virus replication. *World J. Gastroenterol.* 13: 48-64.
6. Liu, X.H., Lin, J., Zhang, S.H., Zhang, S.M., Feitelson, M.A., Gao, H.J. and Zhu, M.H. 2008. COOH-terminal deletion of HBx gene is a frequent event in HBV-associated hepatocellular carcinoma. *World J. Gastroenterol.* 14: 1346-1352.
7. Raimondo, G., Navarra, G., Mondello, S., Costantino, L., Colloredo, G., Cucinotta, E., Di Vita, G., Scisca, C., Squadrito, G. and Pollicino, T. 2008. Occult hepatitis B virus in liver tissue of individuals without hepatic disease. *J. Hepatol.* 48: 743-746.

## SOURCE

Hep B Pol (9F9) is a mouse monoclonal antibody raised against recombinant Hepatitis B Polymerase.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Hep B Pol (9F9) is recommended for detection of amino acids 800-832 corresponding to the RNase H domain of Hepatitis B polymerase by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of Hep B Pol: 95 kDa.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.



See **Hep B Pol (2C8): sc-81590** for Hep B Pol antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.