


AR (N-20): sc-816



The Power to Question

BACKGROUND

Androgens exhibit a wide range of effects on the development, maintenance and regulation of male phenotype and male reproductive physiology. The androgen receptor (AR) is a member of the steroid superfamily of ligand-dependent transcription factors. ARs bind the two biologically active androgens, testosterone (T) and dihydrotestosterone (DHT), with high and nearly identical affinities; however, the rates of association and dissociation of T are about three times more rapid than those of DHT. This difference has resulted in speculation as to whether these differences in binding kinetics could account for the different physiological effects of T and DHT. A striking feature of AR is its rapid degradation in the absence of ligand. It is now well established that androgen binding results in an at least six-fold increase in androgen stability and that ligand-induced stabilization of AR is highly androgen-specific.

CHROMOSOMAL LOCATION

Genetic locus: AR (human) mapping to Xq12; Ar (mouse) mapping to X C3.

SOURCE

AR (N-20) is available as either an affinity purified rabbit (sc-816) or goat (sc-816-G) polyclonal antibody raised against a peptide mapping at the N-terminus of AR of human origin.

PRODUCT

Each vial contains 200 µg/ml IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-816 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-816 X, 200 µg/0.1 ml.

APPLICATIONS

AR (N-20) is recommended for detection of AR of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

AR (N-20) is also recommended for detection of AR in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for AR siRNA (h): sc-29204, AR siRNA (m): sc-29203, AR shRNA Plasmid (h): sc-29204-SH, AR shRNA Plasmid (m): sc-29203-SH, AR shRNA (h) Lentiviral Particles: sc-29204-V and AR shRNA (m) Lentiviral Particles: sc-29203-V.

AR (N-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of AR: 110 kDa.

Positive Controls: T-47D cell lysate: sc-2293 or ZR-75-1 cell lysate: sc-2241.


STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



Western blot analysis of AR expression in AT-3 (A), T-47D (B), ZR-75-1 (C,F) and MCF7 (D) whole cell lysates and SK-BR-3 nuclear extract (E). Antibodies tested include AR (N-20): sc-816 (A-E) and AR (441): sc-7305 (F).



AR (N-20): sc-816. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human prostate carcinoma tissue showing nuclear staining of glandular epithelia (A). Immunofluorescence staining of methanol-fixed LNCaP cells showing nuclear localization (B).

SELECT PRODUCT CITATIONS

- Chen, C.D., et al. 2002. NFκB activates prostate-specific antigen expression and is upregulated in androgen-independent prostate cancer. *Mol. Cell. Biol.* 22: 2862-2870.
- Metzger, E., et al. 2003. A novel inducible transactivation domain in the androgen receptor: implications for PRK in prostate cancer. *EMBO J.* 22: 270-280.
- Chen, Z., et al. 2005. Crucial role of p53-dependent cellular senescence in suppression of Pten-deficient tumorigenesis. *Nature* 436: 725-730.
- Singh, R., et al. 2006. Testosterone inhibits adipogenic differentiation in 3T3-L1 cells: nuclear translocation of androgen receptor complex with β-catenin and T-cell factor 4 may bypass canonical Wnt signaling to down-regulate adipogenic transcription factors. *Endocrinology* 147: 141-154.
- Wissmann, M., et al. 2007. Cooperative demethylation by JMJD2C and LSD1 promotes androgen receptor-dependent gene expression. *Nat. Cell Biol.* 9: 347-353.
- Miyamoto, J., et al. 2007. The pituitary function of androgen receptor constitutes a glucocorticoid production circuit. *Mol. Cell. Biol.* 27: 4807-4814.
- Silva, E.J., et al. 2010. Glucocorticoid receptor in the rat epididymis: expression, cellular distribution and regulation by steroid hormones. *Mol. Cell. Endocrinol.* 325: 64-77.
- Barboro, P., et al. 2011. Androgen receptor and heterogeneous nuclear ribonucleoprotein K colocalize in the nucleoplasm and are modulated by bicalutamide and 4-hydroxy-tamoxifen in prostatic cancer cell lines. *Prostate* 71: 1466-1479.
- Xiao, X., et al. 2011. c-Yes regulates cell adhesion at the blood-testis barrier and the apical ectoplasmic specialization in the seminiferous epithelium of rat testes. *Int. J. Biochem. Cell Biol.* 43: 651-665.

RESEARCH USE

For research use only, not for use in diagnostic procedures.