

Six3 (J-21): sc-81985

BACKGROUND

The Six proteins (sine oculis) are a family of homeodomain transcription factors that share a conserved DNA binding domain. Six3 is required for the specification and proliferation of the eye field in vertebrates and may be involved in some developmental disorders of the brain. Expression of Six3 is detected in human embryos as early as 5-7 weeks of gestation; expression is maintained in the eye throughout the entire period of fetal development. At 20 weeks of gestation, expression of Six3 in the human retina has been observed in ganglion cells and in cells of the inner nuclear layer. Six3 maps to human chromosome 2p21, between genetic markers D2S119 and D2S288. The map position of human Six3 overlaps the positions of two dominant disorders (holoprosencephaly type 2 and malattia leventinese) with ocular phenotypes that have been assigned to this chromosomal region.

REFERENCES

- Granadino, B., et al. 1999. Genomic cloning, structure, expression pattern and chromosomal location of the human Six3 gene. *Genomics* 55: 100-105.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603714. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Del Bene, F., et al. 2004. Direct interaction of Geminin and Six3 in eye development. *Nature* 427: 745-749.
- Hisaoka, M., et al. 2004. Coexpression of NOR-1 and Six3 proteins in extraskeletal myxoid chondrosarcomas without detectable NR4A3 fusion genes. *Cancer Genet. Cytogenet.* 152: 101-107.
- Aijaz, S., et al. 2005. Expression analysis of Six3 and Six6 in human tissues reveals differences in expression and a novel correlation between the expression of Six3 and the genes encoding isocitrate dehydrogenase and cadherin 18. *Genomics* 86: 86-99.
- Conte, I., et al. 2005. Comparative analysis of Six3 and Six6 distribution in the developing and adult mouse brain. *Dev. Dyn.* 234: 718-725.
- Gestri, G., et al. 2005. Six3 functions in anterior neural plate specification by promoting cell proliferation and inhibiting Bmp4 expression. *Development* 132: 2401-2413.
- Lengler, J., et al. 2005. Agonistic and antagonistic action of AP2, Msx2, Pax-6, Prox1 and Six3 in the regulation of Sox2 expression. *Ophthalmic Res.* 37: 301-309.
- Pasquier, L., et al. 2005. First occurrence of aprosencephaly/atelencephaly and holoprosencephaly in a family with a Six3 gene mutation and phenotype/genotype correlation in our series of Six3 mutations. *J. Med. Genet.* 42: e4.

CHROMOSOMAL LOCATION

Genetic locus: SIX3 (human) mapping to 2p21.

SOURCE

Six3 (J-21) is a mouse monoclonal antibody raised against recombinant Six3 of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Six3 (J-21) is recommended for detection of Six3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Six3 siRNA (h): sc-38788, Six3 shRNA Plasmid (h): sc-38788-SH and Six3 shRNA (h) Lentiviral Particles: sc-38788-V.


Molecular Weight of Six3: 37 kDa.

Positive Controls: human thyroid extract: sc-363782.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



Six3 (J-21): sc-81985. Western blot analysis of Six3 expression in human thyroid (diffuse hyperplasia) tissue extract.



Six3 (J-21): sc-81985. Western blot analysis of Six3 expression in 293 whole cell lysate.

SELECT PRODUCT CITATIONS

- Liu, S., et al. 2020. TRIM27 acts as an oncogene and regulates cell proliferation and metastasis in non-small cell lung cancer through SIX3-β-catenin signaling. *Aging*. E-published.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.