

PANK2 (E-17): sc-82288

BACKGROUND

PANK2 (pantothenate kinase 2), also known as HSS, HARP, PKAN or NBIA1, is a ubiquitously expressed 570 amino acid member of the pantothenate kinase family of enzymes that are involved in the synthesis of coenzyme A (CoA). Localized to the cytoplasm and mitochondria, PANK2 is thought to be the chief regulator of CoA biosynthesis, catalyzing the first of five steps in the biosynthetic pathway. Regulated by feedback inhibition from synthesized CoA, PANK2 catalyzes the ATP-dependent conversion of pantothenate to 4'-phosphopantothenate, thus initiating the first committed step in CoA biosynthesis. Defects in the gene encoding PANK2 are the cause of pantothenate kinase-associated neurodegeneration (PKAN) and hypoprebetalipoproteinemia, acanthocytosis, retinitis pigmentosa and pallidal degeneration (HARP). PKAN and HARP are rare disorders characterized by extrapyramidal dysfunction and progressive dementia, both of which are caused by an accumulation of iron in the brain. PANK2 is expressed as three isoforms due to alternative splicing events.

CHROMOSOMAL LOCATION

Genetic locus: PANK2 (human) mapping to 20p13.

SOURCE

PANK2 (E-17) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of PANK2 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82288 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PANK2 (E-17) is recommended for detection of PANK2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with isoform PANK2-2.

Suitable for use as control antibody for PANK2 siRNA (h): sc-76042, PANK2 shRNA Plasmid (h): sc-76042-SH and PANK2 shRNA (h) Lentiviral Particles: sc-76042-V.

Molecular Weight of PANK2 isoform 1: 63 kDa.

Molecular Weight of PANK2 isoform 2: 59 kDa.


Molecular Weight of PANK2 isoform 3: 48 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227 or human liver extract: sc-363766.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



PANK2 (E-17): sc-82288. Western blot analysis of PANK2 expression in Hep G2 whole cell lysate (A) and human liver tissue extract (B).



PANK2 (E-17): sc-82288. Immunoperoxidase staining of formalin fixed, paraffin-embedded human upper stomach tissue showing cytoplasmic staining of glandular cells.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **PANK1/2 (C-4): sc-390595**, our highly recommended monoclonal alternative to PANK2 (E-17).