



XCR1 (S-13): sc-82426

BACKGROUND

G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. G protein-coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. XCR1, also known as GPR5 (G protein-coupled receptor 5) or CCXCR1 (chemokine (C motif) receptor 1), is a 333 amino acid multi-pass membrane protein that belongs to the G protein-coupled receptor 1 family. Functioning as a receptor for select chemokines, XCR1 transduces cellular signals by increasing intracellular calcium ion levels.

REFERENCES

- Larhammar, D., Blomqvist, A.G. and Wahlestedt, C. 1993. The receptor revolution—multiplicity of G-protein-coupled receptors. *Drug Des. Discov.* 9: 179-188.
- Gao, J.L., Kuhns, D.B., Tiffany, H.L., McDermott, D., Li, X., Francke, U. and Murphy, P.M. 1993. Structure and functional expression of the human macrophage inflammatory protein 1 α /RANTES receptor. *J. Exp. Med.* 177: 1421-1427.
- Heiber, M., Docherty, J.M., Shah, G., Nguyen, T., Cheng, R., Heng, H.H., Marchese, A., Tsui, L.C., Shi, X. and George, S.R. 1995. Isolation of three novel human genes encoding G protein-coupled receptors. *DNA Cell Biol.* 14: 25-35.
- Yoshida, T., Imai, T., Kakizaki, M., Nishimura, M., Takagi, S. and Yoshie, O. 1998. Identification of single C motif-1/lymphotactin receptor XCR1. *J. Biol. Chem.* 273: 16551-16554.
- Ji, T.H., Grossmann, M. and Ji, I. 1998. G protein-coupled receptors. I. Diversity of receptor-ligand interactions. *J. Biol. Chem.* 273: 17299-17302.
- Maho, A., Bensimon, A., Vassart, G. and Parmentier, M. 1999. Mapping of the CCXCR1, CX3CR1, CCBP2 and CCR9 genes to the CCR cluster within the 3p21.3 region of the human genome. *Cytogenet. Cell Genet.* 87: 265-268.
- Lee, D.K., Nguyen, T., Lynch, K.R., Cheng, R., Vanti, W.B., Arkhitko, O., Lewis, T., Evans, J.F., George, S.R. and O'Dowd, B.F. 2001. Discovery and mapping of ten novel G protein-coupled receptor genes. *Gene* 275: 83-91.
- Shinkai, H., Morozumi, T., Toki, D., Eguchi-Ogawa, T., Muneta, Y., Awata, T. and Uenishi, H. 2005. Genomic structure of eight porcine chemokine receptors and intergene sharing of an exon between CCR1 and XCR1. *Gene* 349: 55-66.
- Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 600552. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: XCR1 (human) mapping to 3p21.31.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

XCR1 (S-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal extracellular domain of XCR1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82426 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

XCR1 (S-13) is recommended for detection of XCR1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for XCR1 siRNA (h): sc-76931, XCR1 shRNA Plasmid (h): sc-76931-SH and XCR1 shRNA (h) Lentiviral Particles: sc-76931-V.

Molecular Weight of XCR1: 39 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.