



## BAP29 (I-16): sc-82524

### BACKGROUND

BAP29 (BCR-associated protein 29), also known as BCAP29 (B cell receptor-associated protein 29), is a multi-pass membrane protein localizing to the endoplasmic reticulum (ER) and belonging to the BCAP29/BCAP31 family of proteins. It is ubiquitously expressed with predominant expression in brain and testes. BAP29 contains a hydrophobic N-terminus, three transmembrane domains, a coiled-coil region and a C-terminal double-lysine motif that is implicated in vesicular transport. BAP29 exists as a homodimer or as a heterodimer with BAP31 and plays a role in membrane IgD molecule retention in the ER. In addition, the BAP29/BAP31 complex functions as a cargo receptor for MHC class I molecules and is important for recruiting the class I molecules to exit sites of the ER. The BAP29/BAP31 complex is also essential for proper trafficking from the ER to the Golgi.

### REFERENCES

- Kim, K.M., Adachi, T., Nielsen, P.J., Terashima, M., Lamers, M.C., Köhler, G. and Reth, M. 1994. Two new proteins preferentially associated with membrane immunoglobulin D. *EMBO J.* 13: 3793-3800.
- Adachi, T., Schamel, W.W., Kim, K.M., Watanabe, T., Becker, B., Nielsen, P.J. and Reth, M. 1996. The specificity of association of the IgD molecule with the accessory proteins BAP31/BAP29 lies in the IgD transmembrane sequence. *EMBO J.* 15: 1534-1541.
- Breckenridge, D.G., Nguyen, M., Kuppig, S., Reth, M. and Shore, G.C. 2002. The procaspase-8 isoform, procaspase-8L, recruited to the BAP31 complex at the endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* 99: 4331-4336.
- Schamel, W.W., Kuppig, S., Becker, B., Gimborn, K., Hauri, H.P. and Reth, M. 2003. A high-molecular-weight complex of membrane proteins BAP29/BAP31 is involved in the retention of membrane-bound IgD in the endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* 100: 9861-9866.
- Paquet, M.E., Cohen-Doyle, M., Shore, G.C. and Williams, D.B. 2004. BAP29/31 influences the intracellular traffic of MHC class I molecules. *J. Immunol.* 172: 7548-7555.
- Ladasky, J.J., Boyle, S., Seth, M., Li, H., Pentcheva, T., Abe, F., Steinberg, S.J. and Edidin, M. 2006. BAP31 enhances the endoplasmic reticulum export and quality control of human class I MHC molecules. *J. Immunol.* 177: 6172-6181.
- Szczesna-Skorupa, E. and Kemper, B. 2006. BAP31 is involved in the retention of cytochrome P450 2C2 in the endoplasmic reticulum. *J. Biol. Chem.* 281: 4142-4148.
- Zhang, Y. and Williams, D.B. 2006. Assembly of MHC class I molecules within the endoplasmic reticulum. *Immunol. Res.* 35: 151-162.
- Rao, P.S., Bickel, U., Srivenugopal, K.S. and Rao, U.S. 2008. BAP29varP, a variant of BAP29, influences the cell surface expression of the human P-glycoprotein. *Int. J. Oncol.* 32: 135-144.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### CHROMOSOMAL LOCATION

Genetic locus: Bcap29 (mouse) mapping to 12 A3.

### SOURCE

BAP29 (I-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of BAP29 of mouse origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82524 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

BAP29 (I-16) is recommended for detection of BAP29 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for BAP29 siRNA (m): sc-72610, BAP29 shRNA Plasmid (m): sc-72610-SH and BAP29 shRNA (m) Lentiviral Particles: sc-72610-V.

Molecular Weight of BAP29: 29 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.