

FREM2 (S-18): sc-82601

BACKGROUND

FREM2 (FRAS1 related extracellular matrix protein 2) is a 3,169 amino acid single-pass type I membrane protein that localizes to the extracellular side of the cell membrane and contains five Calx- β domains, as well as 12 CSPG repeats. Functioning as an extracellular matrix protein, FREM2 is required for the maintenance of skin and renal epithelia and is also thought to be involved in epidermal adhesion events. Defects or mutations in the gene encoding FREM2, which maps to human chromosome 13, are associated with Fraser syndrome, a multisystem malformation that is characterized by ear abnormalities, congenital heart defects and cutaneous syndactyly. FREM2 exists as multiple alternatively spliced isoforms.

REFERENCES

1. Smyth, I., Du, X., Taylor, M.S., Justice, M.J., Beutler, B. and Jackson, I.J. 2004. The extracellular matrix gene FREM1 is essential for the normal adhesion of the embryonic epidermis. *Proc. Natl. Acad. Sci. USA* 101: 13560-13565.
2. Jadeja, S., Smyth, I., Pitera, J.E., Taylor, M.S., van Haelst, M., Bentley, E., McGregor, L., Hopkins, J., Chalepakis, G., Philip, N., Perez Aytes, A., Watt, F.M., Darling, S.M., Jackson, I., Woolf, A.S. and Scambler, P.J. 2005. Identification of a new gene mutated in Fraser syndrome and mouse myelencephalic blebs. *Nat. Genet.* 37: 520-525.
3. Timmer, J.R., Mak, T.W., Manova, K., Anderson, K.V. and Niswander, L. 2005. Tissue morphogenesis and vascular stability require the FREM2 protein, product of the mouse myelencephalic blebs gene. *Proc. Natl. Acad. Sci. USA* 102: 11746-11750.

CHROMOSOMAL LOCATION

Genetic locus: FREM2 (human) mapping to 13q13.3; Frem2 (mouse) mapping to 3 C.

SOURCE

FREM2 (S-18) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an extracellular domain of FREM2 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82601 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

FREM2 (S-18) is recommended for detection of FREM2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with family members FREM1 or FREM3.

Suitable for use as control antibody for FREM2 siRNA (h): sc-75061, FREM2 siRNA (m): sc-75062, FREM2 shRNA Plasmid (h): sc-75061-SH, FREM2 shRNA Plasmid (m): sc-75062-SH, FREM2 shRNA (h) Lentiviral Particles: sc-75061-V and FREM2 shRNA (m) Lentiviral Particles: sc-75062-V.


Molecular Weight of FREM2: 220 kDa.

Positive Controls: C2C12 Whole Cell Lysate : sc-364188, rat skeletal extract: sc-364810 or mouse embryo extract: sc-364239.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



FREM2 (S-18): sc-82601. Western blot analysis of FREM2 expression in C2C12 (A) and NIH/3T3 (B) whole cell lysates and mouse embryo tissue extract (C).



FREM2 (S-18): sc-82601. Western blot analysis of FREM2 expression in rat skeletal (A) and mouse embryo (B) tissue extracts.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
Satisfation
Guaranteed

Try **FREM2 (F-1): sc-376555**, our highly recommended monoclonal alternative to FREM2 (S-18).