



# Protein C (D-18): sc-82710

## BACKGROUND

Protein C is a vitamin K-dependent plasma protein that is produced in the liver and made up of two polypeptide chains. It is an important anti-coagulant activated by Thrombin bound to an endothelial surface receptor and it enzymatically cleaves activated forms of Factors V and VIII, thus inhibiting blood coagulation. Protein C is similar to the prothrombin group of blood coagulation factors in its primary structure. Normal Protein C concentration in human plasma is approximately 1-3 ng/ml and the proenzyme concentration is approximately 3 µg/ml. Protein C deficiency is associated with inherited thrombophilia, a rare genetic disorder that predisposes affected individuals to venous thrombosis and habitual abortion.

## REFERENCES

- Griffin, J.H., Evatt, B., Zimmerman, T.S., Kleiss, A.J. and Wideman, C. 1982. Deficiency of Protein C in congenital thrombotic disease. *J. Clin. Invest.* 68: 1370-1373.
- Esmon, C.T. 1989. The roles of Protein C and thrombomodulin in the regulation of blood coagulation. *J. Biol. Chem.* 264: 4743-4746.
- Dahlbäck, B., Carlsson, M. and Svensson, P.J. 1993. Familial thrombophilia due to a previously unrecognized mechanism characterized by poor anti-coagulant response to activated Protein C: prediction of a cofactor to activated Protein C. *Proc. Nat. Acad. Sci. USA* 90: 1004-1008.
- Bertina, R.M., Koeleman, B.P., Koster, T., Rosendaal, F.R., Dirven, R.J., de Ronde, H., van der Velden, P.A. and Reitsma, P.H. 1994. Mutation in blood coagulation Factor V associated with resistance to activated Protein C. *Nature* 369: 64-67.
- Dahlbäck, B. 1995. Inherited thrombophilia: resistance to activated Protein C as a pathogenic factor of venous thromboembolism. *Blood* 85: 607-614.
- Preston, F.E., Rosendaal, F.R., Walker, I.D., Briët, E., Berntorp, E., Conard, J., Fontcuberta, J., Makris, M., Mariani, G., Noteboom, W., Pabinger, I., Legnani, C., Scharer, I., Schulman, S. and van der Meer, F.J. 1996. Increased fetal loss in women with heritable thrombophilia. *Lancet* 348: 913-916.
- Abu-Amero, K.K., Owaidah, T.M. and Al-Mahed, M. 2006. Severe type I Protein C deficiency with neonatal purpura fulminans due to a novel homozygous mutation in exon 6 of the Protein C gene. *J. Thromb. Haemost.* 4: 1152-1153.
- Brueckmann, M., Nahrup, A.S., Lang, S., Bertsch, T., Fukudome, K., Liebe, V., Kaden, J.J., Hoffmann, U., Borggreffe, M. and Huhle, G. 2006. Recombinant human activated Protein C upregulates the release of soluble fractalkine from human endothelial cells. *Br. J. Haematol.* 133: 550-557.
- Esmon, C.T. 2006. Inflammation and the activated Protein C anticoagulant pathway. *Semin. Thromb. Hemost.* 1: 49-60.

## CHROMOSOMAL LOCATION

Genetic locus: Proc (mouse) mapping to 18 B1.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## SOURCE

Protein C (D-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Protein C of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82710 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Protein C (D-18) is recommended for detection of Protein C of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Protein C siRNA (h): sc-72054, Protein C shRNA Plasmid (h): sc-72054-SH and Protein C shRNA (h) Lentiviral Particles: sc-72054-V.

Molecular Weight of Protein C: 52 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.