

Rabenosyn-5 (N-20): sc-82730

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. Rabenosyn-5, also known as ZFYVE20 (zinc finger, FYVE domain containing 20), is a 784 amino acid cell membrane protein that contains a C₂H₂-type zinc finger, a FYVE-type zinc finger and a ubiquitin-interacting motif (UIM repeat). The FYVE domain is a cysteine-rich domain of about 70 amino acids. Its primary role is to target signal-transducing proteins to cell membranes through binding to the membrane lipid PIP3 (phosphatidylinositol-3-phosphate) with high specificity. Considered an effector protein, Rabenosyn-5 is required for endosome fusion either homotypically or with clathrin coated vesicles and is involved in the lysosomal trafficking of cathepsin D from the Golgi to lysosomes. Rabenosyn-5 promotes the recycling of transferrin directly from early endosomes to the plasma membrane and binds phospholipid vesicles containing PIP3.

REFERENCES

- Nielsen, E., et al. 2000. Rabenosyn-5, a novel Rab 5 effector, is complexed with hVPS45 and recruited to endosomes through a FYVE finger domain. *J. Cell Biol.* 151: 601-612.
- Hu, Y., et al. 2002. SARA, a FYVE domain protein, affects Rab 5-mediated endocytosis. *J. Cell Sci.* 115: 4755-4763.
- de Renzis, S., et al. 2002. Divalent Rab effectors regulate the sub-compartmental organization and sorting of early endosomes. *Nat. Cell Biol.* 4: 124-133.
- Naslavsky, N., et al. 2004. Rabenosyn-5 and EHD1 interact and sequentially regulate protein recycling to the plasma membrane. *Mol. Biol. Cell* 15: 2410-2422.
- Eathiraj, S., et al. 2005. Structural basis of family-wide Rab GTPase recognition by Rabenosyn-5. *Nature* 436: 415-419.
- Hayakawa, A., et al. 2007. Evolutionarily conserved structural and functional roles of the FYVE domain. *Biochem. Soc. Symp.* 95-105.
- Gengyo-Ando, K., et al. 2007. The SM protein VPS-45 is required for Rab 5-dependent endocytic transport in *Caenorhabditis elegans*. *EMBO Rep.* 8: 152-157.
- Fichtman, B., et al. 2008. EHDS are serine phosphoproteins: EHD1 phosphorylation is enhanced by serum stimulation. *Cell. Mol. Biol. Lett.* 13: 632-648.
- Naslavsky, N., et al. 2009. EHD3 regulates early-endosome-to-Golgi transport and preserves Golgi morphology. *J. Cell Sci.* 122: 389-400.

CHROMOSOMAL LOCATION

Genetic locus: ZFYVE20 (human) mapping to 3p25.1; Zfyve20 (mouse) mapping to 6 D1.

SOURCE

Rabenosyn-5 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Rabenosyn-5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82730 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Rabenosyn-5 (N-20) is recommended for detection of Rabenosyn-5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Rabenosyn-5 (N-20) is also recommended for detection of Rabenosyn-5 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Rabenosyn-5 siRNA (h): sc-76333, Rabenosyn-5 siRNA (m): sc-76334, Rabenosyn-5 shRNA Plasmid (h): sc-76333-SH, Rabenosyn-5 shRNA Plasmid (m): sc-76334-SH, Rabenosyn-5 shRNA (h) Lentiviral Particles: sc-76333-V and Rabenosyn-5 shRNA (m) Lentiviral Particles: sc-76334-V.

Molecular Weight of Rabenosyn-5: 120 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.