

DDX32 (T-20): sc-82859

BACKGROUND

DEAD-box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp, are putative RNA helicases implicated in several cellular processes involving modifications of RNA secondary structure and ribosome/spliceosome assembly. Based on their distribution patterns, some members of this family may be involved in embryogenesis, spermatogenesis, and cellular growth and division. DDX32 (DEAH box polypeptide 32), also known as DHLP1 or DHX32, is a 743 amino acid nuclear protein that localizes to the mitochondria and is a member of the DEAD box helicase family. Expressed in various tissues, DDX32 is up-regulated by Ionomycin in T lymphocytes and down-regulated in acute lymphoblastic leukemia. Considered a novel RNA helicase, DDX32 may play an important role in the development of colorectal cancer and may be involved in regulating T-cell response to certain apoptotic stimuli.

REFERENCES

1. Abdelhaleem, M. 2002. The novel helicase homologue DDX32 is down-regulated in acute lymphoblastic leukemia. *Leuk. Res.* 26: 945-954.
2. Abdelhaleem, M., Sun, T.H. and Ho, M. 2005. DHX32 expression suggests a role in lymphocyte differentiation. *Anticancer Res.* 25: 2645-2648.
3. Alli, Z., Nam, E.H., Beimnet, K. and Abdelhaleem, M. 2005. The activation-induced expression of DHX32 in Jurkat T cells is specific and involves calcium and nuclear factor of activated T cells. *Cell. Immunol.* 237: 141-146.
4. Alli, Z., Ho, M. and Abdelhaleem, M. 2005. Expression of DHX32 in lymphoid tissues. *Exp. Mol. Pathol.* 79: 219-223.
5. Alli, Z., Ackerley, C., Chen, Y., Al-Saud, B. and Abdelhaleem, M. 2006. Nuclear and mitochondrial localization of the putative RNA helicase DHX32. *Exp. Mol. Pathol.* 81: 245-248.
6. Alli, Z., Chen, Y., Abdul Wajid, S., Al-Saud, B. and Abdelhaleem, M. 2007. A role for DHX32 in regulating T-cell apoptosis. *Anticancer Res.* 27: 373-377.
7. Chen, Y., Alli, Z., Ackerley, C., Al-Saud, B. and Abdelhaleem, M. 2007. Altered distribution of heat shock protein 60 (HSP 60) with dysregulated expression of DHX32. *Exp. Mol. Pathol.* 82: 256-261.
8. Alli, Z., Ho, M., Ackerley, C. and Abdelhaleem, M. 2007. Characterization of murine DHX32. *Exp. Mol. Pathol.* 83: 115-118.
9. Huang, C., Liang, X., Huang, R. and Zhang, Z. 2009. Upregulation and clinical relevance of novel helicase homologue DHX32 in colorectal cancer. *J. Exp. Clin. Cancer Res.* 28: 11.

CHROMOSOMAL LOCATION

Genetic locus: DHX32 (human) mapping to 10q26.2; Dhx32 (mouse) mapping to 7 F3.

SOURCE

DDX32 (T-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of DDX32 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-82859 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DDX32 (T-20) is recommended for detection of DDX32 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

DDX32 (T-20) is also recommended for detection of DDX32 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for DDX32 siRNA (h): sc-77143, DDX32 siRNA (m): sc-77144, DDX32 shRNA Plasmid (h): sc-77143-SH, DDX32 shRNA Plasmid (m): sc-77144-SH, DDX32 shRNA (h) Lentiviral Particles: sc-77143-V and DDX32 shRNA (m) Lentiviral Particles: sc-77144-V.

Molecular Weight of DDX32: 84 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **DDX32 (H-3): sc-398743**, our highly recommended monoclonal alternative to DDX32 (T-20).